ABSTRACT

THE VITAMIN A NUTRITION OF PREPARTUM HEREFORD HEIFERS FED DRY LAND RATIONS WITH OR WITHOUT SUPPLEMENTS OF

ALFALFA OR FISH LIVER OIL

Submitted by Dean B. Selleck

In partial fulfillment of the requirements for the Degree of Master of Science Colorado Agricultural and Mechanical College Fort Collins, Colorado

S-2-18-09-03-049

949

5

U18400 9074925

LIERARY COLORADO A. & M. COLLEGE FORT COLLINS, COLDRAND

#### ABSTRACT

Dean B. Selleck Fort Collins, Colorado May 27, 1949 Master of Science

Is vitamin A supplement necessary for the growth of beef heifers under dryland feeding conditions? If vitamin A supplement is needed, can fish liver oil be replaced by small amounts of alfalfa, which is grown extensively in Colorado? What results, as judged by gross observation, may be obtained through moderate vitamin A supplementation in the conditioning of heifers for future reproduction? To secure information pertinent to the above questions, this research was undertaken.

The effect of vitamin A deficiency in the bovine has been studied extensively. No information relative to vitamin A nutrition in preparation of eastern Colorado range heifers for reproduction is available, however. Although many and varied symptoms due to vitamin A deficiency appear, the diagnosis, especially in the mild forms of deficiency, may be difficult. Hart, Mead and Guilbert, 1933, found the degree of deficiency varied in the same herd due to differences in previous feed supply and the amount of reserves stored by the animals. The onset of vitamin A deficiency symptoms was also influenced by the stage of maturity, gestation, production, and rate of growth. Keener, Bechdel, Guerrant and Thorp, 1942, reported that the carotene requirement for calves during severe winter weather was more than twice as great as during warm weather.

Various figures, dependent upon different criteria of evaluation, have been reported for the carotene and vitamin A requirements of cattle. Guilbert, Howell and Hart, 1940, stated that 953 to 1226 I. U. of vitamin A or 1180 to 1500 micrograms of carotene per 100 pounds live weight was the minimum daily requirement for beef cattle. For the growing calf, a ration containing 3000 micrograms carotene per 100 pounds was reported as satisfactory by Boyer et al., 1942; and Moore, Berry and Sykes, 1943. Lewis and Wilson, 1945, were able to maintain growth in calves on 32 U.S.P. units of vitamin A daily per kilogram of body weight. Maximum growth was obtained on an intake of 64 U.S.P. units per kilogram. Blood level and liver storage were low, however, in all calves receiving up to 128 units. From the standpoint of both maximum growth and liver storage, the daily requirement of vitamin A of young calves was considered to be about 250 U.S.P. units, or eight times the minimum requirement.

Gestation presents an added nutritional demand. The requirement of a pregnant heifer for vitamin A is greater than that of a nonpregnant female. The pregnant heifer stores many substances beyond the requirements for herself and fetus. If the need for these reserves fail to develop, the excess is

excreted after calving. Water, nitrogen and vitamin A are well known examples of such storage. Because of this phenomenon, it is difficult to classify the diet of any pregnant female within narrow limits or to make dogmatic statements concerning minor deviations. This yet unmeasured, increased demand of the pregnant female has provided the opportunity for unwarranted advertising and sales claims. According to Kuhlmann and Gallup, 1940, and Moore, 1944, 4000 to 5000 micrograms per 100 pounds body weight was the daily amount of carotene necessary for reproduction, while Davis and Madsen, 1941, stated that a daily intake of 2.72 milligrams of carotene per 100 pounds body weight was sufficient previous to and throughout the gestation period. Recent work at the Beltsville station has apparently confirmed these findings.

Forty Hereford heifer calves were allotted with respect to weight, type, condition, and color, and were fed the experimental rations shown in Table 1.

Lot No.	I	II	III	IV
Winter Rations	Chopped cane Soybean meal Mineral mix Salt	Chopped cane Soybean meal Mineral mix Salt Vit. A oil	Chopped cane Soybean meal Mineral mix Salt Alfalfa hay	Soybean meal Mineral mix Salt Native pasture

Table 1. -- ALLOTTING AND FEEDING PLAN

Summer During the summer of 1947, 1 to 4 heifers Rations in each lot were periodically interchanged between crested wheatgrass and native pasture near Sterling, Colorado. All other heifers were pastured on native pasture. Moorman's mineral mixture was fed.

> During the summer of 1948, all heifers were pastured on native pasture near Idalia, Colorado, and bonemeal was used as the mineral supplement.

During the wintering periods, all lots received concentrates once and roughages twice daily. All feeds and refuse were weighed each day. Feed samples taken every 28 days throughout the two years of the experiment, were reduced to an air-dry basis in a large drier, then ground in a Wiley mill and preserved for chemical analysis in air-tight glass jars.

The vitamin A oil was mixed with the protein supplement daily and fed in a manner to insure equal distribution to all animals in the lot. Alfalfa hay was fed to Lot III at a level to provide carotene equivalent in vitamin A value to that received as oil concentrate by the animals of Lot II. Vitamin A equivalent intake was calculated by dividing the carotene intake by the factor 3.5.

Dry matter consumption by heifers on winter range was calculated according to Morrison's recommendations. Dry matter consumption on summer range was computed from values obtained for steers on pasture by Garrigus and Rusk.

Two day weights of each heifer were obtained at the beginning and end of each phase, and individual single day weights were taken each 28 days. All animals were graded for type at the beginning of each phase, and for condition at the beginning of the experiment and the end of the second wintering period. In grading the heifers for both type and condition, individuals were given the regular market grades from common minus to choice plus, then the grades were assigned a numerical value of 0 to 10 in order to determine the average of the lot. All heifers were number branded during the spring of 1947, and were then clipped at frequent intervals to facilitate identification and the keeping of individual records.

Blood samples were taken from the jugular vein of each animal at intervals not exceeding 56 days. The samples were refrigerated until analyses could be made, and care was taken to analyze as quickly as possible. Serum carotene and vitamin A were determined colorimetrically by the Kimble method, using an Aminco type F photoelectric colorimeter. A factor was applied in the computation of vitamin A to correct for the blue color developed by carotene. Serum calcium was determined by the Tisdall method as modified by Clark and Collip, and serum phosphorus was measured by a modification of the Bell-Doisy phosphate method as described by Briggs.

The vitamin A content of the oil fed to Lot II was determined colorimetrically by the Carr-Price method as modified by Dann and Evelyn. Feed samples were analyzed for dry matter, total carotenoids and total nitrogen. Carotenoids were determined by the method described by Peterson, and nitrogen analysis was carried out by the Kjeldahl method. All analyses were conducted in the nutrition laboratories in the Animal Investigations Section at Colorado A & M College.

In the absence of classical vitamin A deficiency symptoms, together with evidence of growth and parturition performance currently considered as normal in practical beef herd maintenance, it may be concluded that common dryland rations, of the quality observed during this experiment, support adequate vitamin A nutrition. Within the limitation of this study, one may also conclude that alfalfa hay is equal to, and in some cases superior to, fish liver oil as a vitamin A supplement in the maintenance of serum carotene and vitamin A levels, weight gains, type, and condition grades, vigor, and general appearance.

Limited observations indicate that an advantage was derived from the feeding of alfalfa hay supplement in that calving time for the lot was appreciably shorter than that of the other lots, being completed in the interval of one estrus cycle of 21 days. Seventy per cent of the calves of this lot were dropped in nine days. Furthermore, the calves dropped by the alfalfa supplemented lot were by far the most vigorous of all.

Lot	I	II	III	IV
	Control	Fish liver oil	Alfalfa	Pasture
(Winter (Summer <u>Total Gains</u> (Winter (Summer (Overal Initial weight (12)	$\begin{array}{ccccccc} 1946-47 & 119.75 \\ 1947 & 147.80 \\ 1947-48 & 197.45 \\ 1948 & 114.00 \\ 1 & 579.00 \\ -12-46) & 303.50 \end{array}$	127.75 156.45 200.80 117.50 602.50 305.25	131.25 160.00 200.00 103.75 595.00 304.25	104.75 190.66 124.06 180.83 600.30 292.75
Final weight (11-6 Serum carotene (Maximu Mcgms/100 ml. ( (Minimu	$ \begin{array}{cccc} -48) & 882.50 \\ m & 637.35 \\ (7-8-48) \\ m & 39.72 \\ (5-6-48) \end{array} $	907.75 576.90 (7-8-48) 35.88 (5-6-48)	899.25 610.65 (7-8-48) 41.40 (5-6-48)	892.77 874.83 (7-8-48) 68.70 (12-12-46)
Serum vitamin A (Maximu I.U./100 ml. ( (Minimu	$\begin{array}{cccc} m & & 438.96 \\ & & (11-5-48) \\ m & & 100.87 \\ & & (4-2-47) \end{array}$	418.62 (1-8-48) 115.77 (4-2-47)	527.07 (6-12-47) 118.77 (4-2-47)	579.23 (5-6-48) 156.81 (4-2-47)
Vitamin A equivalent (Maximu Intake (Units/100# Da.)(	n $45,177$ (5-15 to 6-12-47)	44,604 (5-15 to 6-12-47)	44,155 (5-15 to 6-12-47)	45,984 (5-15 to 6-12-47)
(Minimu	m 857 (4-8 to 5-6-48)	873 (9-30 to 11-5-48)	878 (9-30 to 11-5-48)	328 (2-5-to 3-4-48)
Days required for 70% of calve in lot to be dropped	s 25	80 10	100 9	100 17
Days required for complete cal Average birth weights (pounds) Cows requiring assistance at c	ving 56 72.6 alving 2	35 72.1 2	21 71.9 0	49 68.6 0

Table 2.--COMPOSITE DATA OF VITAMIN A NUTRITION AND REPRODUCTION

THESIS

-----

THE VITAMIN A NUTRITION OF PREPARTUM HEREFORD HEIFERS FED DRY LAND RATIONS WITH OR WITHOUT SUPPLEMENTS OF ALFALFA OR FISH LIVER OIL

Submitted by Dean B. Selleck

In partial fulfillment of the requirements for the Degree of Master of Science Colorado Agricultural and Mechanical College

Agricultural and Mechanical College Fort Collins, Colorado

May, 1949

COLORADO A & M. COLLEGE

COLORADO AGRICULTURAL AND MECHANICAL COLLEGE May 28, 194 9 WE HEREBY RECOMMEND THAT THE THESIS PREPARED UNDER OUR SUPERVISION BY DEAN B. SELLECK ENTITLED THE VITAMIN A NUTRITION OF PREPARTUM HEREFORD HEIFERS FED DRY LAND RATIONS WITH OR WITHOUT SUPPLEMENTS OF ALFALFA OR FISH LIVER OIL BE ACCEPTED AS FULFILLING THIS PART OF THE REQUIREMENTS FOR THE DEGREE OF MASTER OF SCIENCE. CREDITS 15 Committee on Graduate Work 2. 2. 4 ast Minor Professor Major Professor of Department Dean of Division Committee on Final Examination Examination Satisfactory LE, Wash a ney Dean of the Graduate School

2

### ACKNOWLEDGMENT

The author wishes to express his appreciation for the cooperation, interest and technical advice received from Dr. S. S. Wheeler, Head of the Animal Husbandry Department, Dr. L. E. Washburn, Professor of Animal Nutrition and Professor W. E. Connell during the course of this research.

To Mrs. John Nishamura and Mr. Marvin E. Kniese for their assistance in analysis of the many blood and feed samples, to Miss Hazel Brandis for typing and proofing the manuscript, and to members of the Akron Dryland Experiment Station and ranchers who fed and cared for the cattle during the wintering and summering periods, the author is grateful.

## TABLE OF CONTENTS

Chapter							Page
I	INTRODUCTION			•			7
II	REVIEW OF LITERATURE			•			9
III	METHODS AND MATERIALS				• •		18
IV	ANALYSIS OF DATA						21
V	General observations Eye disorders						21 22 23 26 26 28 30 32 34
	General observations Eye disorders Serum carotene and vitamin A Serum calcium and phosphorus Weight gains Feed consumption Cost of live weight gains Type and condition Calving performance	• • • • • • • • •	• • • • • • • •	•	• •	• • • • • • • • •	34 34 35 37 38 39 40
VI	SUMMARY			•	• •		41
	APPENDIX		•	•	• •		43
	BIBLIOGRAPHY			•	•		124

LIST OF TABLES

Tabl	e	Page
1	ALLOTTING AND FEEDING PLAN	18
2	FEED CONSUMPTION AND WEIGHT GAINS	29
3	FEED COST PER 100 POUNDS GAIN	30
4	SUMMARY OF CALVING DATA	33

## LIST OF FIGURES

Figure	2	Page
l	RELATIONSHIP OF VITAMIN A EQUIVALENT INTAKE AND BLOOD SERUM LEVELS (LOTS I AND IV)	24
2	RELATIONSHIP OF VITAMIN A EQUIVALENT INTAKE AND BLOOD SERUM LEVELS (LOTS II AND III)	25
3	AVERAGE WEIGHTS OF HEIFERS BY LOTS	27
4	TYPE AND CONDITION GRADE	31

# Chapter I INTRODUCTION

Is vitamin A supplement necessary for the growth of beef heifers under dryland feeding conditions? If vitamin A supplement is needed, can fish liver oil be replaced by small amounts of alfalfa, which is grown extensively in Colorado? What results, as judged by gross observation, may be obtained through moderate vitamin A supplementation in the conditioning of heifers for future reproduction? To secure information pertinent to the above questions, the research reported in this thesis was undertaken.

Previous work has shown that at times it may be necessary to supply beef cattle with a vitamin A supplement. Following analysis of several hundred samples of winter forage, which comprises a major part of the annual feed consumed by beef cattle, Washburn (55) stated, "Many animals in Colorado may suffer from prolonged "borderline" carotene intake, resulting in reduced resistance to infections, impairment of growth, reproduction and lactation." Based on this analysis, calculations indicated that it was impossible for cattle to consume sufficient amounts of some poor quality sorghum forages, alfalfa, and native hays and most of the winter range grass to meet minimum requirements for vitamin A.

8

The importance of beef production as an agricultural enterprise in Colorado emphasizes the need for finding a practical, economical, and efficient method of supplementing vitamin A if and when such is needed.

# Chapter II REVIEW OF LITERATURE

0

The effect of vitamin A deficiency in the bovine has been studied extensively. These studies reveal that the carotene and vitamin A requirements may be altered by many conditions such as growth, reproduction, lactation, source of carotene, weather, previous feed conditions, age, sex, and breed.

A blood plasma carotene level of 25 micrograms per 100 milliliters was reported as adequate for health of Hereford and Shorthorn heifers by Davis and Madsen (13). Earlier, Moore (40) found that the plasma carotene of Holstein and Ayrshire calves should be 0.2 micrograms per milliliter or above to maintain fair general health and to prevent nyctalopia. These findings are in agreement with those of Kuhlman and Gallup (35) who obtained satisfactory growth responses in Jersey calves with plasma carotene values ranging from 25 to 100 micrograms per 100 cc. Boyer et al. (1) stated that blood plasma vitamin A is a more delicate measure of the state of vitamin A nutrition than is either growth or blood carotene, and that a blood plasma vitamin A level of 10 micrograms or more per 100 cc. is necessary for adequate vitamin A nutrition of the growing calf. Blood plasma vitamin A levels below 7-8 micrograms per 100 cc. are definitely inadequate. This work showed that the blood plasma carotene levels of 50-70 micrograms per 100 cc. for Holsteins and 110-140 micrograms per 100 cc. for Guernseys are necessary to maintain adequate blood vitamin A. Hansen and associates (23) confirmed these values by showing that the newborn calf has a very low average blood plasma vitamin A level, and that normal concentrations of the order of 9-10 micrograms per 100 milliliters or more must be quickly attained (1-2 days) for a favorable chance of survival.

Blood plasma carotene levels of 78-96 micrograms per 100 milliliters are required for normal reproduction in Hereford and Shorthorn heifers, according to Davis and Madsen (13) whereas, Kuhlman and Gallup (35) and Gallup and Kuhlman (17) reported that 150 micrograms were necessary for Jersey cows and heifers. Payne and Kingman (48) found that the blood plasma carotene level of first calf range Hereford heifers must be at least  $117.75 \stackrel{+}{-} 7.21$ micrograms per 100 ml. to support normal gestation. The requirement for aged cows was  $82.88 \stackrel{+}{-} 4.11$  micrograms per 100 ml. Kuhlman and Gallup (33) also reported that the carotene requirements for first calf heifers may be somewhat higher than those for older cows, as the younger animals need carotene for growth in addition to that demanded for maintenance and reproduction.

Many authors have reported carotene requirements on the basis of body weight. Guilbert and Hart (19) published data stating that the minimum daily carotene requirement of the bovine was 26-33 micrograms per kilogram of body weight. Later work by Guilbert, Howell and Hart (21), and Guilbert, Miller and Hughes (22) confirmed the above carotene requirement and set the minimum vitamin A requirement at 5.1-6.4 micrograms per kilogram of body weight. Converse and Meigs (10) started four day old calves on a low vitamin A basal ration supplemented with varying amounts of carotene (29-87 micrograms per kilogram of body weight) in oil or with cod liver oil supplying 7.55-22.6 micrograms per kilogram. Calves at all levels of intake died or developed subnormally. Lewis and Wilson (36) were able to maintain fair growth in calves on 32 U.S.P. units per kilogram of body weight. From the standpoint of both normal growth and liver storage, the daily intake of vitamin A of young calves should be about eight times the minimum requirement. For the growing calf a ration containing 3000 micrograms of carotene per 100 pounds per day was reported as satisfactory by Boyer et al. (1) and Moore, Berry and Sykes (45). Ward, Bechdel and Guerrant (54) were able to maintain growth and prevent usual vitamin A deficiency symptoms in dairy calves by supplying a ration containing 11 micrograms of carotene per pound of body weight.

These authors also stated that the adequacy of carotene level depended largely upon the source of supply.

Gestation and lactation produces an added nutritional demand, and the requirement of a pregnant heifer for vitamin A is greater than that of a similarly healthy but nonpregnant female (11). Converse and Meigs (9) found that when timothy or clover hay was fed without pasture a daily intake of 80 to 130 milligrams of carotene was sufficient to enable production of normal calves, but that weak calves were the rule when the daily intake of carotene was 50 to 60 milligrams. These values are higher than those suggested by Kuhlman and Gallup (34), and Moore (43), in which cases 40 to 50 micrograms per pound of body weight were given as the minimum amount of carotene necessary for either reproduction or lactation. Davis and Madsen (13) fed 60 micrograms of carotene per kilogram to heifers and reported the birth of apparently normal calves. These results should not be considered as conflicting, since the breeds of animals used, sources of carotene and other conditions of the experiments were not comparable.

Investigations by Keener et al. (29) and by Kempt (30) indicated that the vitamin A and carotene requirements of dairy calves were only one-half as great during the summer as during the winter.

The development of symptoms of avitaminosis A depends largely upon the previous feed conditions and the

length of time elapsing prior to depletion of the vitamin A reserves. Vitamin A deficiency in cattle under natural conditions was reported by Hart, Mead and Guilbert (26) who found that during the very dry summer of 1932, cows suffered from night blindness, corneal ulceration, diarrhea, and nonrecurrence of estrus after calving. Calves born late in the season died, and the oil from as much as 10 grams of their livers gave no blue color in the Carr-Price test. Survivors recovered when green feed was again available. Fernandes (15) also found that drought and the lack of green fodder resulted in vitamin A deficiency as evidenced by general debility and blindness among cattle. Feeding of green fodder and cod liver oil yielded good results.

Jones and co-workers (28), (50) stated that in seven experiments conducted from 1934 to 1941 with 310 feeder cattle ranging from 3 to 16 months of age at the time of being placed in the feedlot and fed rations low in carotene, the time required for depletion of body reserves of vitamin A, as indicated by night blindness, ranged from 45 to 268 days. Reserves of young animals became depleted in less time than that of older animals. In dry years less time was required for depletion than in years with favorable rainfall. Earlier work by Dickson and Jones (14) indicated that steer calves would suffer from vitamin A deficiency in approximately 100 days, but these steers would regain good health following the addition of one pound of high quality alfalfa hay to the daily ration. Other investigators have published similar findings. Schmidt (52) working with feedlot cattle in the Red Beds region of Texas, found that cows and calves fed largely on white kafir, cottonseed meal and cottonseed hulls developed sore eyes, nasal discharge, marked weight losses, and convulsions after approximately 120 days of feeding. These conditions were corrected by adding alfalfa, cod liver oil, or crystalline carotene supplements to the ration. Guilbert and Hart (20) reported clinical symptoms of vitamin A deficiency occurring in beef steers after 225 days on a ration of dried molasses beet pulp, rolled barley, cottonseed meal, and calcium carbonate. Reserves of vitamin A had been practically exhausted after 282 days, and the animals were in a critical condition.

Many authors have reported on variations in the carotenoid content of forage plants. In 1932 Converse and Meigs (8) observed that cows receiving overripe timothy hay as the only source of roughage for a 12-month period failed to produce normal calves. Hart and Guilbert (24) reported reproductive failure in range cattle due to vitamin A deficiency appearing near the end of a dry feed season. Hart, Guilbert and Goss (25) analyzed over 400 samples of range forage collected during different seasons over a two year period. Marked changes in nutritive value were evident during various stages of growth and in different seasons. A very high rate of mortality of newborn calves occurred in a herd of 250 Holstein cows pastured during the winter months in corn stalk fields and then fed in a dry lot on a ration of wheat straw and 12 to 14 bushels of yellow corn per day (51). Flora and co-workers (16) fed growing calves poor quality, bleached, and aged timothy hays with a grain mixture. The resulting avitaminosis A was corrected quite effectively by feeding carrots, dehydrated alfalfa hay, or by a commercial carotene concentrate.

Following a study of seasonal changes of the blood carotenoid and vitamin A levels, and the normal ratio of these levels in cattle, Braun (3) concluded that seasonal changes of carotenoid intake were unaffected by pathological disturbances, but varied with statistical significance according to the age and breed of the animals. Seasonal changes in the vitamin A level were dependent on carotene and vitamin A intake, but were modified by parturition, abortion, and acute infections. The vitamin Acarotene ratio was below normal during parturition, abortion and acute infections, but was slightly above normal if vitamin A supplement was fed. Data collected by Sutton and Soldner (53) indicated wide fluctuations in blood plasma carotene within the different breeds responding similarly to changes in carotene intake. Blood plasma vitamin A was found to vary between narrower limits than

did blood plasma carotene. Changes in blood plasma vitamin A tended to lag behind the blood plasma carotene changes. In another publication, Braun (4) stated that, "The utilization of stored vitamin first forces available carotenoid stores to be converted to vitamin A, thus decreasing the carotenoid level without decreasing the vitamin A level." A tendency towards a direct relationship between vitamin A stores and the vitamin A level of the blood was found to exist only when the former fell below normal levels.

Findings of Nelson and associates (47) revealed that the carotene and vitamin A content of blood plasma was higher in beef calves than in dairy calves. Moore and Berry (44) found the plasma vitamin A of Holstein, Ayrshire, and Guernsey calves from birth to four months of age varied from 7.2 to 14 micrograms per cent. The feeding of a vitamin A supplement was recommended where difficulty is encountered in raising calves. Moore (42) showed that within a breed heifers had definitely lower plasma carotene values than producing cows, under both winter and summer pasture conditions. He described the rise of plasma carotene from 0.10 to 0.33 micrograms per cc. at 15 hours and to 8.49 micrograms per cc. at 168 hours following the feeding of freshly cut alfalfa to a cow with a very low initial plasma carotene value.

Many symptoms have been attributed to carotene and vitamin A deficiencies. Moore (41) found that calves receiving low carotene rations developed nyctalopia, papillary edema, and a permanent type of blindness due to a constriction of the optic nerve. Symptoms of vitamin A deficiency were associated with generalized edema or anasarca in cattle by Madsen and associates (37). Crowther (11) reported the occurrence of irregular estrus, probable prolonged gestation, and difficult parturition. The placenta was often abnormal and fetal death and resorption or abortion frequently resulted in cases of vitamin A deficiency.

Bulls subjected to a period of vitamin A depletion showed a marked loss of fertility (11). Sexual activity and ability decreased rapidly during the period of depletion, and semen samples from vitamin A deficient bulls showed marked increases of cellular debris and abnormal spermatozoa with a decline in motility (38).

## Chapter III

## METHODS AND MATERIALS

Forty Hereford heifer calves were allotted with respect to weight, type, condition, and color, and were fed the experimental rations shown in Table 1.

Table 1.--ALLOTTING AND FEEDING PLAN

Lot No.	I	II	III	IV	
Winter Rations	Chopped cane Soybean meal Mineral mix Salt	Chopped Chopped Chopped cane cane cane Soybean Soybean Soybean meal meal meal Mineral Mineral Mineral mix mix mix Salt Salt Salt Vit. A Alfalfa oil hay		Soybean meal Mineral mix Salt Native pasture	
Summer Rations	During the summer of 1947, 1 to 4 heifers in each lot were periodically interchanged between crested wheatgrass and native pasture near Sterling, Colorado. All other heifers were pastured on native pasture. Moorman's mineral mixture was fed.				
	During the pastured o rado, and supplement	summer of 19 n native past bonemeal was	948, all heife Sure near Idal used as the m	ers were Lia, Colo- Dineral	
concentrate	During the w	intering peri roughages twi	lods, all lots .ce daily. Al	received	

and refuse were weighed each day. Feed samples taken every 28 days throughout the two years of the experiment were reduced to an air-dry basis in a large drier, then ground in a Wiley mill and preserved for chemical analysis in airtight glass jars.

The vitamin A oil was mixed with the protein supplement daily and fed in a manner to insure equal distribution to all animals in the lot. Alfalfa hay was fed to Lot III at a level to provide carotene equivalent in vitamin A value to that received as oil concentrate by the animals of Lot II. Vitamin A equivalent intake was calculated by dividing the carotene intake by the factor 3.5.

Dry matter consumption by heifers on winter range was calculated according to Morrison's recommendations (46). Dry matter consumption on summer range was computed from values obtained for steers on pasture by Garrigus and Rusk (18).

Two day weights of each heifer were obtained at the beginning and end of each phase, and individual single day weights were taken each 28 days. All animals were graded for type at the beginning of each phase, and for condition at the beginning of the experiment and the end of the second wintering period. In grading the heifers for both type and condition, individuals were given the regular market grades from common minus to choice plus then the grades were assigned a numerical value of 0 to 10 in order to determine the average of the lot. All heifers were number branded during the spring of 1947, and were then clipped at frequent intervals to facilitate identification and the keeping of individual records.

Blood samples were taken from the jugular vein of each animal at intervals not exceeding 56 days. The samples were refrigerated until analyses could be made, and care was taken to analyze as quickly as possible. Serum carotene and vitamin A were determined colorimetrically by the Kimble method (31) using an Aminco type F photoelectric colorimeter. A factor was applied in the computation of vitamin A to correct for the blue color developed by carotene (2). Serum calcium was determined by the Tisdall method as modified by Clark and Collip (7), and serum phosphorus was measured by a modification of the Bell-Doisy phosphate method as described by Briggs (5).

The vitamin A content of the oil fed to Lot II was determined colorimetrically by the Carr-Price method as modified by Dann and Evelyn (12). Feed samples were analyzed for dry matter, total carotenoids and total nitrogen. Carotenoids were determined by the method described by Peterson (49), and nitrogen analysis was carried out by the Kjeldahl method. All analyses were conducted in the nutrition laboratories of the Animal Investigations Section at Colorado A & M College.

# Chapter IV ANALYSIS OF DATA

### General observations

Data relative to weight, condition scores, and blood analysis generally reflected the influence of feed and environmental conditions. The health of all lots was excellent and only very slight differences, if any, could be attributed to diseased, parasitic, or injured animals. Death loss was limited to one heifer from the pasture lot, caused by poisoning while on summer range. Slight infestations of lice were encountered at times, but were corrected by spraying or dusting before the condition became serious. Considerable shrink resulted from shipping the heifers to and from summer pasture. Change of feed usually enabled the animals to regain quickly the weight lost long before the next weigh day.

Classical symptoms of vitamin A deficiency were not observed during the course of the experiment. Appearance of such symptoms were no doubt prevented by the excellent condition of the pastures and the high quality of the cane and alfalfa, resulting from above average annual rainfall. The heifers which received a small amount of alfalfa showed more bloom, scored higher in type and condition, showed more vigor, and had better appetites than the other groups. Animals receiving continuous pasture appeared rough and thin after the winter months. These animals made, however, the largest gains while on summer pasture resulting in an appearance equal to the other lots by the end of the summer.

### Eye disorders

Some cases of eye disorder diagnosed as "pink eye" were encountered soon after the animals were placed on summer pasture in 1947. Six heifers were affected, one each from the control lot, oil supplemented lot, and pasture lot, and three from the alfalfa supplemented lot. Heifers from the control, oil supplemented and pasture lots were not affected to the point of blindness and recovered in from four to six days. In the alfalfa supplemented lot, animal number 15 was blind in one eye for three weeks, and animals number 27 and 31 were blind in one or both eyes for about six weeks. During this period the carotene ranged from 500 to 600 micrograms and the vitamin A varied from 527 to 1120 I.U. per 100 ml. of serum.

### <u>Serum</u> carotene

Lot averages of blood serum carotene ranged from a low of 36 micrograms per 100 milliliters in the fish liver oil lot for May of 1948 to a high of 875 micrograms

in the pasture lot for July 1948. The alfalfa supplemented lot maintained higher serum carotene values than did either the control or fish liver oil supplemented lots. Changes in carotene consumption were reflected by parallel changes in serum carotene as shown in Figures 1 and 2. The fish liver oil lot showed lower serum carotene levels than the control lot which received the same carotene intake. Some animals were found repeatedly to have higher individual carotene values than others under the same conditions receiving the same feed.

### Serum vitamin A

Serum vitamin A reached a low of 101 I.U. per 100 milliliters in April 1947, in the control lot. The highest averages occurred in May of 1948 when the pasture lot reached a peak of 579 I.U. per 100 milliliters of serum. Serum vitamin A values were not as variable as the carotene values. Changes in serum vitamin A tended to lag behind changes in serum carotene. A noticeable decline in serum vitamin A occurred in all lots during the summer months although serum carotene remained high. Figures 1 and 2 show for all lots the vitamin A values compared with the vitamin A equivalent intake.



Fig. 1.--Relationship of vitamin A equivalent intake and blood serum levels





### Serum calcium and phosphorus

Values for serum calcium and phosphorus were variable, ranging from 7.06 to 12.84 milligrams of calcium and from 4.48 to 10.38 milligrams of phosphorus per 100 milliliters of serum. No one lot had consistently higher or lower levels than other lots. Phosphorus values declined and calcium values increased with the approach of maturity. In this experiment the concentration of Ca and PO<sub>4</sub> ions in the blood serum were generally in reciprocal relation to each other. An increase in the concentration of one resulted in a decrease in the concentration of the other. Although the mineral consumption by the pasture lot was much lower than by the other lots, there were, however, no appreciable differences in the calcium and phosphorus content of the serum.

### Weight gains

Average weights for each lot of animals for each month are shown in Figure 3. As calves and yearlings, slightly heavier weights were attained by the lot receiving alfalfa while the fish liver oil lot weighed heaviest at two years of age. Both vitamin A and carotene supplemented lots showed nearly equal weights and gains superior to those of the check lot throughout the experiment. The pasture lot made the largest gains during the summer periods.

1000 400 600 800 0 DEC, 12 1946 □ @< APR. 30 1947 - • • • • • (CX .... -SEPT, 28 1947 OR 690 . DATE COx. DO ON a . FEB. 5 1948 X...LOT 3 D...LOT 4 ... LOT I O... LOT 2 OK ox JUNE 26 1948 . ..... .... 000 O MO NOV. 11 1948 000

LIVE WEIGHT (LBS.)
#### Feed consumption

Data in Table 2 show the average daily dry matter, protein, carotene, and vitamin A equivalent intake for each lot and experimental period. Dry matter, protein, and carotene fed were equal for lots I and II. Differences shown on the table are due to feed refused and weighed back, and to differences in live weights between the lots. Calculations for feed consumption on pasture were based on values for dry matter intake as published by Morrison (46) or by Garrigus and Rusk (18).

#### Type and condition

Figure 4 shows graphically the average type and condition scores given by different judges to the lots at various periods of the experiment. At the beginning of the experiment, all lots were judged as equal in type and practically equal in condition. As time passed, the alfalfa supplemented lot gained the superior rating in both type and condition scores. The fish liver oil supplemented lot was surpassed by the control lot and was approached closely by the pasture lot in both type and condition score.

Tabl	e 2FEED	CONSUMPT	ION AND W	EIGHT GAI	NS	
Lot	Average weight	Average gain	Average dry mat- ter intake 100# live wt./day	Average protein intake pounds 100# live wt./day	Average carotene intake mg/100# live wt./day	Average vitamin A intake *** units/ 100# live wt./day
	Pounds	<u>Pounds</u> Wi	Pounds .nter 1946	Pounds -47	Mgs.	<u>Units</u>
I II III IV	379.32 384.46 385.10 344.76	119.75 127.75 131.25 104.75	2.33 2.22 2.29 2.49*	0.245 0.234 0.269 0.221	9.961 9.089 13.123 20.020	2,846 4,038 3,749 5,720
			Summer 19	47		
I II III IV	521.50 534.04 537.40 529.78	147.80 156.45 160.00 190.66	2.60** 2.55** 2.54** 2.57**	0.186 0.182 0.180 0.181	59.694 58.600 58.494 59.268	17,055 16,743 16,713 16,934
		Wi	nter 1947	-48		
I II III IV	692.66 712.12 717.00 662.54	197.45 200.80 200.00 124.06	2.37 2.32 2.28 1.99*	0.175 0.170 0.174 0.128	5.134 5.011 5,568 5,252	1,467 2,374 1,591 1,500
			Summer 19	48		16426
I II III IV	817.85 848.39 843.10 823.28	114.00 117.50 103.75 180.83	1.83** 1.79** 1.80** 1.83**	0.083 0.081 0.082 0.083	11.556 11.307 11.345 11.536	3,302 3,231 3,242 3,296
*	Dry matter was calcul	consumpt ated by u	ion for a sing Morr	nimals on ison's re	winter r commendat	ange ions.
**	Dry matter obtained f and Rusk. Bulletin 4	consumpt for steers Illinois 54.	ion was co on pastu Agricult	omputed f re publis ural Expe	rom value hed by Ga riment St	s rrigus ation
***	Vitamin A grams carc vitamin A.	equivalen otene equi	nt intake valent to	calculate l Intern	d as 3.5 ational u	micro- nit

Period	Lot I	Lot II	Lot III	Lot IV
Winter 1946-47	\$14.08	\$13.38	\$12.79	\$11.77
Summer 1947	5.07	4.79	4.69	3.93
Winter 1947-48	18.95	18.77	17.68	14.45
Summer 1948	10.53	10.21	11.57	6.64
Overall	12.74	12.33	12.04	8.29

# Cost of live weight gains

Table 3 shows the feed cost per 100 pounds gain for each lot during each phase and for the entire experiment. Overall cost of 100 pounds gain was as follows: Lot IV, \$8.29; Lot III, \$12.04; Lot II, \$12.33, and Lot I, \$12.74. Cost of 100 pounds gain was more expensive in the second winter and summer than during the first winter and summer.

Table 3 .-- FEED COST PER 100 POUNDS GAIN



# Calving performance

Information on parturition assembled in Table 4 reveals that all animals conceived. Four calves were born dead, two each in the control and fish liver oil lots. The time required for conception, as judged by calving dates was: Alfalfa lot, 21 days; fish liver oil lot, 35 days; pasture lot, 49 days; and control lot, 56 days. Two heifers each in the control and fish liver oil lots required assistance during parturition. One case of placental retention occurred in each lot.

Table 4SUMMARY OF CAL	VING DATA			
	Lot I	Lot II	Lot III	Lot IV
Number of cows	10	10	10	9
Number of calves	10	10	10	9
Birth date of first calf	3/10/49	2/9/49*	3/13/49	3/13/49
Birth date of last calf	5/5/49	4/20/49	4/3/49	5/1/49
Days required for 70% of calves in lot to be dropped	25	(10)	9	17
Days required for complet calving	te 56	(35)	21	49
Average birth weight (pounds)	72.6	72.1	71.9	68.6
Death loss of cows during calving	g 1	0	0	0
Death loss of calves	2	2	0	0
Cows requiring assist- ance at calving	2	2	0	0
Retained placentas	l	1	1	1

\* Calf was not carried the full gestation period and was dead at birth. First live calf was born March 16, 1949. Chapter V DISCUSSION

#### General observations

Due to above average rainfall during both summers, superior quality of the summer range and winter forage apparently inhibited the development of gross vitamin A deficiency symptoms. Although such circumstances are the exception rather than the rule in the high plains area, certainly during the time of this experiment the common dryland rations of eastern Colorado satisfied the vitamin A requirements for growth and conception of beef heifers. Nevertheless, the animals receiving a small amount of alfalfa hay exhibited better appetites, more vigor and more bloom to their hair coats than did the others. By the end of the summer periods, differences in the general appearance between lots were not distinguishable.

## Eye disorders

"Pink eye" is rather frequently observed on the northern plains range soon after young animals are placed on spring pasture. In this experiment, those heifers which received higher levels of carotene from alfalfa

during the previous winter were the most severely affected. Most of the animals in the alfalfa lot presented a yellowish lachrymal discharge, although only three developed severe inflammation and blindness. One is inclined to incriminate hypercarotenemia as a cause for this trouble. Certainly, the observed serum carotene and vitamin A levels cannot logically indicate deficiency. A carotenevitamin A-thyroid interrelationship is suggested as a possible contributing factor in "pink eye". It is known that excess carotene depresses the activity of the thyroid gland, and that thyroidectomy is frequently followed closely by "pink eye".

#### Serum carotene and vitamin A

Blood serum carotene of animals receiving alfalfa hay was considerably higher than that of the other lots during the wintering periods. Although extreme variation was observed in the pasture lot, the serum carotene promptly increased rather rapidly from the lowest observed values with the consumption of new grass early in the spring, and attained the highest level during the summer periods. Kniese (32) pointed out the possibility of low levels acting as a stimulus to the rate of conversion by stating, "....indicates an increased ability to convert carotene from plant sources to serum carotene."

The serum carotene of the fish liver oil lot was usually lower than that of the control lot which received the same carotene intake. This indicates that fish liver oil depresses serum carotene in some manner not explained by the data obtained in this work.

At the same vitamin A equivalent consumption, alfalfa supplementation resulted in higher serum vitamin A values than obtained from fish liver oil supplement. Only in a few instances, when the fish oil lot received temporarily higher vitamin A equivalent intake, did the serum vitamin A values approach those recorded for the alfalfa supplemented animals. Higher serum vitamin A levels were obtained during the second winter although the vitamin A equivalent intake per 100 pounds of body weight was lower than during the previous winter. The available data does not permit an explanation as to whether this was due to greater storage of vitamin A during the previous summer, lower requirements with increased age, or because of gestation. An appreciable decline in serum vitamin A occurred in all lots during the summer months, while at the same time the serum carotene remained high. It would appear logical to explain this phenomena as being due to the depressing effect of hot weather upon the thyroid gland (6) and the fact that carotene cannot be converted into vitamin A during hypothyroidism (39) or in the absence of the thyroid gland (6).

#### Serum calcium and phosphorus

Mineral mixture was self-fed to all lots and

variation with respect to consumption in the different lots might partly explain the considerable variation in the serum calcium and phosphorus values. No logical explanation can be given for a low mineral consumption by the pasture lot during the winter months, or for the maintenance of normal serum values throughout this period of diminished mineral intake. In general, the concentration of Ca and  $PO_4$  ions in the blood serum were in reciprocal relationship to each other. An increase in the concentration of the one resulted in a decrease in the concentration of the other. The serum calcium values tended to increase and the serum phosphorus levels decrease with the approach of maturity.

#### Weight gains

Throughout the greater part of the experiment the alfalfa supplemented lot was the heaviest, although the fish oil lot weighed the most near the end of the second summer. Difference in weight of the two lots was so slight at all times, however, that it is impossible from weight data alone to critically evaluate either supplement. Although no classical symptoms of vitamin A deficiency were in evidence, there were temporary periods during which it was impossible, due to the quality of the feed, to supply carotene intakes considered as minimum for growth by some authors. According to other recommendations however, the carotene consumption at these times was ade-

quate. During these periods the vitamin A equivalent intake was greater in the fish liver oil and alfalfa supplemented lots. This might account for the superior gains of the vitamin A supplemented lots. The small gains made by the pasture lot during the winters were followed by superior gains during the summer, resulting in total body weights equal to the vitamin A supplemented lots toward the end of the summer periods.

#### Feed consumption

The heifers receiving alfalfa exhibited the best appetites as shown by the small amount of feed refused and weighed back. Stimulated appetites may be due either to some factor in the alfalfa hay, or merely to increased palatability brought about by the variety introduced into the ration. Dry matter consumption for the groups in drylot was about 2.3 pounds daily per 100 pounds of body weight throughout both winters. Poor quality of the feed during the winter of 1947-48 resulted in lower consumption of protein and carotene than the previous winter. Calculated intakes of dry matter, protein, and carotene were considerably higher in the summer of 1947, than in the summer of 1948.

# Cost of live weight gains

Animals which received pasture made the least expensive live weight gain of all lots throughout the

entire experiment. The alfalfa lot showed the least expensive gain of the three drylot fed groups. Gain of the control lot was the most expensive. Total feed cost was highest for the fish liver oil lot, but relatively greater gain reduced the cost per 100 pounds gain. Cost of gain is based largely on economic considerations and obviously does not critically evaluate biologic performance, which in this case concerned vitamin A nutrition. A lower cost of gain as shown by the heifers on continuous pasture does not mean therefore a more adequate vitamin A metabolism. Data pertinent to the gestation and lactation periods of these animals and also the performance of their offspring must first be obtained before final recommendations can be made.

#### Type and condition

All lots were scored as equal with regard to type and approximately equal in condition at the start of the experiment. Although the heifers were graded several times by different judges, the alfalfa lot was rated repeatedly as superior in type. This lot was followed in the order given by the control lot, fish liver oil lot, and the pasture lot. Condition scores varied according to the feed conditions prior to judging. The alfalfa lot was given the highest score for condition, the pasture lot was lowest and the other lots were rated approximately equal. In view of the fact that the alfalfa lot was consistently rated superior in both type and condition, it is logical to assume that the alfalfa supplement was responsible for the attractive appearance of this lot when viewed by the judges.

## Calving performance

Observations limited to behavior of cows and calves at parturition indicate that the period of conception, as judged by calving dates, was shortest in the alfalfa supplemented lot. This lot was followed by the fish liver oil lot, which required 14 days longer time for calving. The time required to drop 70 per cent of the calves in each lot was as follows: Alfalfa supplemented lot, 9 days; fish liver oil supplemented lot, 10 days; pasture lot, 17 days; and control lot, 25 days. Animals receiving the unsupplemented, common dryland rations required considerably longer time for conception after pasture mating with the bull. Heifers receiving continuous pasture dropped calves weighing the least at birth. This might be due to the lower dry matter, protein, or carotene consumption by the dams. Calf birth weight averages in the other lots were practically equal. The loss of one cow from the control lot during parturition was due to an abnormal position of the calf and probably cannot be attributed to nutritional factors.

# Chapter VI SUMMARY

The vitamin A nutrition of prepartum Hereford heifers was studied under practical dry land conditions, supplementing commonly fed rations with alfalfa hay and fish liver oil. Forty animals were allotted to four groups which received the following rations: (a) cane, (b) cane and fish liver oil, (c) cane and alfalfa, and (d) continuous pasture. All lots received in addition soybean oil meal, salt, and mineral mixture. The alfalfa hay and fish liver oil were supplemented on a vitamin A equivalent basis. All groups grazed together on summer range.

Individual two day average weights were taken at the beginning and end of each phase, and single day weights were obtained every 28 days. Blood samples were collected at approximately 56 day intervals, and the serum was analyzed for carotene, vitamin A, calcium, and phosphorus. Samples of all feeds were analyzed every 28 days for dry matter, nitrogen, and carotenoids.

In the absence of classical vitamin A deficiency symptoms, together with evidence of growth and parturition performance currently considered as normal in practical beef herd maintenance, it may be concluded that common dry land rations, of the quality observed during this experiment, support adequate vitamin A nutrition. Within the limitations of this study, one may also conclude that alfalfa hay is equal to, and in some cases superior to, fish liver oil as a vitamin A supplement in the maintenance of serum carotene and vitamin A levels, weight gains, type and condition grades, vigor, and general appearance.

Limited observations indicate that an advantage was derived from the feeding of alfalfa hay supplement in that calving time for the lot was appreciably shorter than that of the other lots, being completed in the interval of one estrus cycle of 21 days. Seventy per cent of the calves of this lot were dropped in nine days. Furthermore, the calves dropped by the alfalfa supplemented lot were by far the most vigorous of all.



LIST OF APPENDIX TABLES

Table		Page
l	TOTAL FEED CONSUMPTION - WINTER 1946-47	46
2	FEED ANALYSIS 1946-48	54
3	TOTAL DRY MATTER, PROTEIN, CAROTENE AND VITAMIN A EQUIVALENT CONSUMPTION - WINTER 1946-47.	57
4	TOTAL DRY MATTER, PROTEIN, CAROTENE AND VITAMIN A EQUIVALENT CONSUMPTION -	-
	SUMMER 1947	61
5	TOTAL DRY MATTER, PROTEIN, CAROTENE AND VITAMIN A EQUIVALENT CONSUMPTION - WINTER 1947-48	65
6	TOTAL DRY MATTER, PROTEIN, CAROTENE AND VITAMIN A EQUIVALENT CONSUMPTION - SUMMER 1948	60
7	PLOOD SEDIM CADOTENE (Norma/100 MI ) 10/6 /7	09
(	BLOOD SEROM CAROTENE (Megms/100 MI.) 1940-47.	13
8	BLOOD SERUM CAROTENE (Mcgms/100 Ml.) 1947-48.	77
9	BLOOD SERUM VITAMIN A (I. U./100 Ml.) 1946-47	81
10	BLOOD SERUM VITAMIN A (I. U./100 Ml.) 1947-48	85
11	BLOOD SERUM PHOSPHORUS (Mgms/100 Ml.) 1946-47	89
12	BLOOD SERUM PHOSPHORUS (Mgms/100 Ml.) 1947-48	93
13	BLOOD SERUM CALCIUM (Mgms/100 Ml.) 1946-47	97
14	BLOOD SERUM CALCIUM (Mgms/100 Ml.) 1947-48	101

		LIST OF	APPENI	DIX TA	BLE	cs(	Cont	in	ue	ed				
Table														Page
15	BODY	WEIGHTS	WINTER	1946-	47	(POUI	NDS)	•		•	•	•	•	105
16	BODY	WEIGHTS	SUMMER	1947	(P(	DUNDS	).	•		•		•	•	109
17	BODY	WEIGHTS	WINTER	1947-	48	(POUI	NDS)	•	•				•	113
18	BODY	WEIGHTS	SUMMER	1948	(PC	DUNDS	).	•		•	•	•	•	117
19	TYPE	AND CONI	DITION (	GRADES	5.			•	•	•		•	•	121
20	FEED	COSTS 19	946-48.											123

0.00 10 0.00 10 0.00 10	10.00 10.00	 			
0.00 10	10.00		2,970.00	12/12-1/8	I
0.00 10	20.00	 	3,360.00	1/8-2/5	
0.00	20.00	 	2,780.00	2/5-3/5	
0.00 20	20.00	 	2,800.00	3/5-4/2	
0.00 20	20.00	 	2,780.00	4/2-4/30	
0.00 10	10.00	 	1,525.00	4/30-5/15	
0.00 80	90.00	 	16,215.00	Total	
0.06 (	0.06	 	10.46	Av/Hd/Day	
	9	 	1,525.00 16,215.00 10.46	4/30-5/15 Total Av/Hd/Day	

Lot No.	Dates	Chopped cane lbs.	Alfalfa hay lbs.	Bleached cane lbs.	Protein supple- ment lbs.	Vitamin A oil ml.	Mineral mixture lbs.	Salt 1bs.
II	12/12-1/8	2,970.00			187.50	120.00	10.00	10.00
	1/8-2/5	3,360.00			210.00		10.00	10.00
	2/5-3/5	1,950.00		900.00	210.00	64.00	10.00	30.00
	3/5-4/2	1,850.00		930.00	210.00	48.00	30.00	20.00
	4/2-4/30	1,850.00		950.00	210.00	48.00	20.00	20.00
	4/30-5/15	1,525.00			108.75	24.00	10.00	10.00
	Total	13,505.00		2,780.00	1,136.25	304.00	90.00	100.00
	Av/Hd/Day	8.71		1.79	0.73	0.20	0.06	0.06

Lot No,	Dates	Chopped cane lbs.	Alfalfa hay lbs.	Bleached cane lbs.	Protein supple- ment lbs.	Vitamin A oil ml.	Mineral mixture lbs.	Salt lbs.
III	12/12-1/8	2,430.00	540.00		129.00		10.00	10.00
	1/8-2/5	2,800.00	560.00		140.00		10.00	10.00
	2/5-3/5	2,240.00	560.00		135.00		20.00	40.00
	3/5-4/2	2,240.00	560.00		140.00		40.00	40.00
	4/2-4/30	2,240.00	560.00		140.00		20.00	20.00
	4/30-5/15	1,230.00	295.00		72.50		10.00	10.00
	Total	13,180.00	3,075.00		756.50		110.00	130.00
	Av/Hd/Day	8.50	1.98		0.49		0.07	0.08

Lot No.	Dates	Chopped cane lbs.	Alfalfa hay lbs.	Bleached cane lbs.	Protein supple- ment lbs.	Vitamin A oil ml.	Mineral mixture lbs.	Salt lbs.
IV	12/12-1/8	480.00			187.50			
	1/8-2/5				210.00			
	2/5-3/5				202.50		10.00	20.00
	3/5-4/2				210.00			
	4/2-4/30				210.00			
	4/30-5/15				108.75			
	Total	480.00			1,128.75		10.00	20.00
	Av/Hd/Day	0.31			0.73		0.006	0.013

A

Lot No.	Dates	Chopped cane lbs.	Alfalfa hay lbs.	Protein supple- ment lbs.	Vitamin A oil ml.	Mineral mixture lbs.	Salt lbs.
I	11/14-12/11	5,840.00		270.00		20.00	30.00
	12/11-1/8	6,150.00		280.00		10.00	20.00
	1/8-2/5	6,160.00		280.00		10.00	10.00
	2/5-3/4	6,160.00		280.00		10.00	20.00
	3/4-4/8	7,630.00		350.00		20.00	30.00
	4/8-5/6	5,366.00		280.00			10.00
	Total	37,306.00		1,740.00		70.00	120.00
	Av/Hd/Day	21.32		0.99		0.04	0.07

Lot No.	Dates .	Chopped cane lbs.	Alfalfa hay lbs.	Protein supple- ment lbs.	Vitamin A oil ml.	Mineral mixture lbs.	Salt lbs.
II	11/14-12/11	5,840.00		270.00	40.00	20.00	30.00
	12/11-1/8	6,150.00		280.00	140.00	10.00	30.00
	1/8-2/5	6,160.00		280.00	140.00	10.00	10.00
	2/5-3/4	6,160.00		280.00	20.00	10.00	20.00
	3/4-4/8	7,667.00		350.00	25.00	20.00	30.00
	4/8-5/6	5,518.00		280.00	20.00	10.00	30.00
	Total	37,495.00		1,740.00	385.00	\$0.00	150.00
	Av/Hd/Day	21.43		0.99	0.22	0.05	0.09

KA

Lot No.	Dates	Chopped cane lbs.	Alfalfa hay lbs.	Protein supple- ment lbs.	Vitamin A oil ml.	Mineral mixture lbs.	Salt lbs.
III	11/14-12/11	5,560.00	270.00	202.50		20.00	30.00
	12/11-1/8	5,871.50	280.00	210.00		10.00	30,00
	1/8-2/5	5,880.00	280.00	210.00		10.00	10.00
	2/5-3/4	5,600.00	560.00	140.00		10.00	20.00
	3/4-4/8	6,974.00	680.00	175.00		20.00	40.00
	4/8-5/6	4,980.00	560.00	140.00			30.00
	Total	34,865.50	2,630.00	1,077.50		70.00	160.00
	Av/Hd/Day	19.92	1.50	0.62		0.04	0.09

Lot No.	Dates	Chopped cane lbs.	Alfalfa hay lbs.	Protein supple- ment lbs.	Vitamin A oil ml.	Mineral mixture lbs.	Salt 1bs.
IV	11/14-12/11			243.00		10.00	10.00
	12/11-1/8			243.00			
	1/8-2/5	720.00		252.00			10.00
	2/5-3/4	360.00		252.00		20.00	20.00
	3/4-4/8	360.00		315.00			20.00
	4/8-5/6			252.00			10.00
	Total	1,440.00		1,557.00		30.00	70.00
	Av/Hd/Day	0.91		0.99		0.02	0.04
							1

Table 2	-FEED ANALY	SIS 1946-	48		
Type of feed	Date of sample	Master number	Percent dry matter	Lbs. protein per lb. D.M.	Mcgms. carotene per 1b. D.M.
Chopped cane	12/12/46 12/23/46 1/8/47 2/5/47 3/5/47 4/2/47 4/2/47 4/30/47 5/14/47 11/14/47 12/11/47 1/8/48 2/5/48 3/4/48 3/4/48 4/8/48 5/6/48	1,361 1,364 1,379 1,396 1,457 B 1,500 1,532 1,558 1,767 1,779 1,779 1,792 1,812 1,844 1,845 1,860 1,874	58.24 49.95 65.78 80.46 88.68 93.52 75.45 93.55 75.99 75.45 75.45 92.62 90.42	.0783 .0740 .0797 .0765 .0716 .0828 .0562 .0759 .0607 .0576 .0741 .0595 .0559 .0261 .0291 .0337	9,128 11,222 4,968 2,170 3,330 4,788 586 4,077 3,884 4,026 1,217 2,525 2,944 1,177 1,372 1,200
Bleached cane	1/8/47 2/5/47 3/5/47	1,378 1,395 1,457 B	73.77 59.72 80.46	.0676 .0666 .0716	1,946 1,915 3,330
Alfalfa hay	12/12/46 12/23/46 1/8/47 2/5/47 3/5/47 4/2/47 4/30/47 5/14/47 12/11/47 1/8/48 2/5/48 3/4/48 4/8/48 5/6/48	1,362 1,365 1,380 1,397 1,456 1,501 1,533 1,559 1,778 1,778 1,791 1,811 1,843 1,859 1,873	88.68 89.93 80.12 88.54 90.08 91.74 88.22 87.62 89.87 86.94 86.65 85.39 91.06 92.95	.1388 .1799 .1862 .1842 .1840 .1888 .2044 .1908 .1771 .1765 .1803 .1946 .1888 .2039	12,414 13,357 11,608 12,346 9,173 9,972 12,178 11,604 3,877 5,108 7,902 4,273 5,121 2,486
Pasture grass	12/12/46 1/8/47 2/5/47 3/5/47	1,360 1,377 1,394 1,458	89.78 91.24 91.30 96.31	.0403 .0349 .0456 .0364	2,296 1,886 1,004 561

Type of feed	Date of sample	Master number	Percent dry matter	Lbs. protein per lb. D.M.	Mcgms. carotene per lb. D.M.
Pasture grass cont.	4/2/47 4/30/47 5/14/47 6/12/47 7/10/47 8/7/47 9/4/47 10/2/47 11/14/47 12/11/47 12/11/47 1/8/48 2/5/48 3/4/48 4/8/48 5/6/48 5/6/48 5/6/48 5/6/48 5/6/48 5/6/48 5/6/48 9/2/48 9/30/48 11/6/48	1,499 1,534 1,560 1,655 1,665 1,6689 1,766 1,768 1,766 1,768 1,766 1,768 1,786 1,786 1,814 1,866 1,877 1,888 1,887 1,888 1,888 1,888 1,888 1,903	93.01 44.11 56.58 37.57 54.51 71.04 75.69 73.11 85.18 52.80 84.80 79.57 78.91 62.41 96.70 82.11 96.70 82.11 96.728 67.14 75.61 88.36 96.08	.0449 .1318 .0782 .1026 .0715 .0521 .0525 .0459 .0339 .0475 .0384 .0294 .0167 .0169 .0494 .0628 .0673 .0518 .0496 .0496 .0440 .0448	790 37,450 24,915 58,959 24,939 13,500 9,926 1,862 15,835 938 567 884 000 1,427 10,727 11,719 6,559 7,093 8,410 3,505 25
Crested wheat- grass	6/12/47 7/10/47 8/7/47 9/4/47 10/2/47 11/14/47	1,654 1,660 1,674 1,688 1,703 1,765	27.91 48.50 66.02 77.03 53.74 51.26	.0919 .0737 .0972 .0502 .1300 .1133	27,814 16,746 25,068 6,391 34,342 31,280
Soybean oil meal	12/12/46 1/8/47 2/5/47 3/5/47 4/2/47 4/20/47 11/14/47 12/11/47	1,357 1,374 1,391 1,478 1,496 1,529 1,764 1,777	90.56 89.84 89.37 91.19 88.77 89.64 90.46 90.66	.4330 .4521 .4754 .4526 .4407 .4352 .4457 .4475	193 000 000 000 000 315 52

Table 2.--FEED ANALYSIS 1946-48--Continued

Type of feed	Date of sample	Master number	Percent dry matter	Lbs. protein per lb. D.M.	Mcgms. carotene per lb. D.M.
Soybean oil meal cont.	1/8/48 2/5/48 3/4/48 4/8/48 5/6/48	1,790 1,810 1,842 1,858 1,872	89.72 90.78 90.74 92.41 92.11	.4678 .4519 .4765 .4432 .4677	137 000 000 54 29
Fish liver oil	8/15/46 8/18/47 2/3/48 6/14/48	Vitamin Vitamin Vitamin Vitamin	A = 26,000 A = 31,000 A = 33,683 A = 33,500	) I. U. pe ) I. U. pe ; I. U. pe ) I. U. pe	r gram r gram r gram r gram

Table 2.--FEED ANALYSIS 1946-48--Continued

		Dry ma	tter	Prot	ein	Caro (Mg	tene ms.)	Vitamin A e	quivalen s)*
Lot No.	Dates	Total	Av.per 100# per da.	Total	Av.per 100# per da.	Total	Av.per 100# per da.	Total	Av.per 100# per da.
I	12/12 to 1/8	1,832	2.09	202.19	0.229	15,208	17.23	4,345,114	4,922
	1/8 to 2/5	2,630	2.69	277.91	0.285	9,375	9.60	2,678,525	2,743
	2/5 to 3/5	2,415	2.27	253.70	0.239	6,720	6.33	1,919,857	1,809
	3/5 to 4/2	2,557	2.28	267.22	0.238	9,612	8.56	2,746,203	2,446
	4/2 to 4/30	2,735	2.32	284.22	0.241	11,293	9.56	3,226,496	2,732
	4/30 to 5/15	1,532	2.40	156.15	0.244	6,360	9.95	1,817,120	2,816
	Average		2.34		0.246		10.20		2,846
	Total	13,701	1	,441.39		58,568		16,733,315	- 5.0

		Dry m	atter	Prot	Protein (lbs.)		tene ms.)	Vitamin A equivalen (units)*	
lo.	Dates	Total	Av.per 100# per da.	Total	Av.per 100# per da.	Total	Av.per 100# per da.	Total	Av.per 100# per da.
I	12/12 to 1/8	1,832	2.06	202.19	0.228	15,208	17.13	7,735,280	8,712
	1/8 to 2/5	2,630	2.68	277.91	0.283	9,375	9.54	2,678,480	2,725
	2/5 to 3/5	2,325	2.17	245.56	0.229	6,411	5.98	3,640,280	3,397
	3/5 to 4/2	2,374	2.09	246.82	0.217	7,547	6.63	3,512,880	3,086
	4/2 to 4/30	2,549	2.12	262.59	0.219	9,263	7.71	4,003,160	3,331
	4/30 to 5/15	1,532	2.34	156.15	0.239	6,360	9.74	2,495,400	3,790
	Average		2.24		0.236		9.46		4,038
	Total	13,242	1	,391.22		54,156		24,065,480	

Г

		Dry m	atter s.)	Prot (1b	Protein (lbs.)		tene ms.)	Vitamin A equivaler (units)*	
ot	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
II	12/12 to 1/8	1,947	2.18	236.05	0.265	18,405	20.66	5,258,705	5,904
	1/8 to 2/5	2,632	2.67	304.32	0.309	13,466	13.68	3,847,423	3,915
	2/5 to 3/5	2,414	2.26	281.98	0.264	10,792	10.09	3,083,383	2,883
	3/5 to 4/2	2,529	2.22	297.37	0.261	12,560	11.04	3,588,687	3,155
	4/2 to 4/30	2,682	2.23	316.85	0.263	14,825	12.31	4,235,813	3,518
	4/30 to 5/15	1,481	2.25	171.34	0.261	8,282	12.61	2,366,265	3,543
	Average		2.30		0.270		13.40		3,749
	Total	13,685	1	,607.91		78,330		22,380,276	

		Dry m (lbs	atter .)	Prot (1b	ein s.)	Caro (Mg	tene ms.)	Vitamin A e (unit	quivalent s)*
Lot No.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
V	12/12 to 1/8	2,240	2.65	147.82	0.175	6,242	7.37	1,783,340	2,106
	1/8 to 2/5	2,296	2.60	185.36	0.210	3,046	3.45	870,303	984
	2/5 to 3/5	2,352	2.55	176.13	0.191	1,696	1.84	484,617	525
	3/5 to 4/2	2,436	2.48	180.55	0.184	1,519	1.55	433,992	443
	4/2 to 4/30	2,548	2.42	370.15	0.292	53,536	50.88	15,295,907	14,512
	4/30 to 5/15	1,410	2.38	185.71	0.314	40,942	69.23	11,697,705	18,597
	Average		2.51		0.228		22.39		5,720
	Total	13,282	1	,182.72		106,981		30,565,864	

		Dry m	atter	Prote (1b;	ein s.)	Caro (Mg	tene ms.)	Vitamin A e (unit	quivalen s)*
.ot Da Io.	ites	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
I 5/15	to 6/12	3,612	3.00	362.86	0.302	190,473	158.12	54,420,919	45,177
6/12	to 7/10	3,682	2.88	317.24	0.248	139,980	109.57	39,994,238	31,307
7/10	to 8/7	3,780	2.67	251.44	0.178	73,927	52.29	21,121,992	14,941
8/7 t	0 9/4	3,850	2.49	250.79	0.163	54,431	35.16	15,551,800	10,046
9/4 t	0 10/2	3,892	2.41	286.98	0.179	61,041	37.82	17,440,384	10,806
10/2	to 10/30	3,892	2.40	203.00	0.124	34,359	21.17	9,816,736	6,220
10/30	) to 11/14	2,085	2.42	87.28	0.108	15,476	17.96	4,421,669	5,131
Avera	ıge		2.61		0.186		61.72		17,661
Total		24,793		1,759.59		569,687		162,767,738	1

		Dry m (lb	atter s.)	Prot (1b	ein s.)	Carot (Mgr	cene ns.)	Vitamin A ed (units	quivalen s)*
Lot No.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
I :	5/15 to 6/12	3,640	2.96	365.67	0.298	191,937	156.15	54,839,200	44,604
6	5/12 to 7/10	3,699	2.84	318.70	0.245	140,614	107.98	40,175,512	30,850
1	7/10 to 8/7	3,794	2.62	252.38	0.174	74,201	51.32	21,200,318	14,664
8	3/7 to 9/4	3,867	2.45	251.89	0.160	54,671	34.70	15,620,243	9,913
	9/4 to 10/2	3,903	2.38	287.81	0.176	61,219	37.35	17,491,015	10,672
	10/2 to 10/30	3,912	2.34	204.05	0.122	34,535	20.66	9,867,173	5,902
2	10/30 to 11/14	2,094	2.34	87.58	0.106	15,517	17.38	4,433,483	4,964
1	Average		2.56		0.183		60.79		17,367
r.	Total	24,909	1	,768.08		572,694		163,626,944	

		Dry m (lb	atter s.)	Prot (1b	ein s.)	Carot (Mgr	cene ns.)	Vitamin A ed (units	quivalent s)*
Lot No.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
III 5	5/15 to 6/12	3,654	2.93	367.18	0.294	192,669	154.54	55,048,340	44,155
6	0/12 to 7/10	3,716	2.81	320.17	0.241	141,268	106.79	40,362,405	30,511
7	7/10 to 8/7	3,797	2.62	252.57	0.174	74,259	51.19	21,216,792	14,627
Ś	1/7 to 9/4	3,867	2.45	251.89	0.160	54,671	34.70	15,620,243	9,914
9	0/4 to 10/2	3,906	2.38	288.04	0.176	61,268	37.32	17,505,128	10,663
l	.0/2 to 10/30	3,914	2.34	204.13	0.122	34,544	20.60	9,869,799	5,885
l	.0/30 to 11/14	2,100	2.33	87.86	0.105	15,573	17.31	4,449,420	4,945
A	verage		2.55		0.182		60.35		17,242
I	otal	24,954	1	,771.84		574,252		164,072,127	
		Dry m (lb	atter s.)	Prot (1b	ein s.)	Carot (Mgr	cene ns.)	Vitamin A ed (units	quivalent s)*
------	--------------	--------------	---------------------------	-------------	---------------------------	---------------	---------------------------	------------------------	---------------------------
lo.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
V 5/	15 to 6/12	3,584	3.05	306.04	0.308	188,978	160.94	53,993,374	45,984
6/	12 to 7/10	3,696	2.85	318.45	0.246	140,508	180.44	40,145,175	30,983
7/	10 to 8/7	3,794	2.62	252.38	0.174	74,201	51.23	21,200,318	14,638
8/	7 to 9/4	3,875	2.45	252.41	0.160	54,785	34.69	15,652,786	9,912
9/	4 to 10/2	3,906	2.37	288.04	0.175	61,268	37.17	17,505,128	10,620
10	/2 to 10/30	3,920	2.33	204.45	0.122	34,600	20.61	9,885,736	5,945
10	/30 to 11/14	1,884	2.35	80.46	0.096	14,552	18.15	4,157,843	5,186
Av	erage		2.57		0.183		71.89		17,609
To	tal	24,659	1	,756.23		568,892		162,540,360	

		Dry m (1b	atter s.)	Prot (1b	ein s.)	Caro (Mg	tene ms.)	Vitamin A e (unit	quivalen s)*
lot Io.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Áv.per 100# per day
I.	11/14 to 12/11	3,833	2.32	321.83	0.194	14,236	8.60	4,067,320	2,457
	12/11 to 1/8	4,610	2.63	402.46	0.230	11,448	6.54	3,270,810	1,867
	1/8 to 2/5	4,917	2.65	427.89	0.231	8,744	4.71	2,498,229	1,347
;	2/5 to 3/4	4,350	2.22	354.25	0.182	11,203	5.72	3,200,731	1,635
	3/4 to 4/8	5,842	2.30	342.32	0.135	9,558	3.76	2,730,886	1,076
1	4/8 to 5/6	5,170	2.45	271.73	0.129	6,328	3.00	1,807,905	857
	Average		2.43		0.184		5.39		1,540
1	Fotal	28,722	2	,120.48		61,517		17,575,881	

		Dry m (lb	atter s.)	Prot (1b	ein s.)	Caro (Mg	tene ms.)	Vitamin A e (unit	quivalen s)*
Lot No.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
I	11/14 to 12/11	3,833	2.25	321.83	0.190	14,236	8.36	5,273,520	3,097
	12/11 to 1/8	4,610	2.55	402.46	0.224	11,448	6.35	7,491,680	4,153
	1/8 to 2/5	4,917	2.57	427.89	0.224	8,744	4.57	6,791,160	3,510
	2/5 to 3/4	4,350	2.16	354.25	0.177	11,203	5.57	3,803,800	1,891
	3/4 to 4/8	5,873	2.26	343.17	0.132	9,598	3.69	3,496,150	1,343
	4/8 to 5/6	5,309	2.45	276.09	0.128	6,506	3.00	2,462,040	1,137
	Average		2.37		0.179		5.26		2,522
1	Total	28,892	2	,125.69		61,735		29,246,350	

		Dry m (lb	atter s.)	Prot (1b	ein s.)	Caro (Mg	tene ms.)	Vitamin A e (unit	quivalen s)*
Lot No.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
II	11/14 to 12/11	3,838	2.23	326.29	0.190	14,618	8.45	4,176,508	2,425
	12/11 to 1/8	4,597	2.52	404.07	0.221	12,040	6.61	3,439,871	1,889
	1/8 to 2/5	4,885	2.53	428.10	0.222	9,924	5.14	2,835,291	1,468
	2/5 to 3/4	4,333	2.14	364.15	0.180	13,120	6.47	3,748,445	1,849
	3/4 to 4/8	5,812	2.23	366.91	0.141	11,545	4.42	3,298,512	1,263
	4/8 to 5/6	5,203	2.39	303.05	0.139	7,827	3.60	2,236,386	1,029
	Average		2.34		0.182		5.78		1,654
	Total	28,668	2	,192.57		69,074		19,735,013	

		Dry m (lb	atter s.)	Prot (1b	ein s.)	Caro (Mg	tene ms.)	Vitamin A e (unit	quivalen s)*
Lot No.	Dates	Total	Av.per 100 # per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
V	11/14 to 12/11	3,100	2.07	222.09	0.148	24,195	16.14	6,912,869	4,612
	12/11 to 1/8	3,175	2.04	200.42	0.129	2,246	1.44	641,844	412
	1/8 to 2/5	3,326	2.01	199.52	0.121	2,890	1.50	825,589	500
	2/5 to 3/4	3,402	2.00	169.38	0.099	1,950	1.15	557,284	328
	3/4 to 4/8	4,252	2.00	265.88	0.124	3,157	1.48	902,089	424
	4/8 to 5/6	3,478	1.99	287.75	0.164	19,736	10.14	5,638,767	2,897
	Average		2.02		0.131		5.31		1,529
	Total	20,733	1	,345.04		54,174		15,478,442	

		Dry m (lb	atter s.)	Prot (1b	ein s.)	Caro (Mg	tene ms.)	Vitamin A economication (units	quivalen 5)*
lot No.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
I	5/6 to 6/5	4,365	1.93	260.15	0.115	39,957	17.65	11,416,500	5,044
	6/5 to 7/8	4,811	1.91	243.94	0.097	34,036	13.50	9,724,440	3,856
	7/8 to 8/5	4,144	1.85	193.94	0.086	30,359	13.53	8,674,120	3,865
	8/5 to 9/2	4,200	1.81	178.08	0.077	32,558	14.01	9,302,400	4,004
	9/2 to 9/30	4,284	1.77	153.80	0.064	25,524	10.54	7,292,592	3,010
	9/30 to 11/5	5,425	1.75	206.15	0.067	9,575	3.09	2,735,750	884
	Average		1.84		0.084		12.05		3,444
	Total	27,229	1	,236,06		172,009		49,145,802	

1 International Unit vitamin A.

		Dry m (lb	atter s.)	Prot (1b	ein	Caro (Mg	tene ms.)	Vitamin A ed	quivalen s)*
lo.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
I	5/6 to 6/5	4,410	1.87	262.84	0.112	40,369	17.14	11,534,100	4,897
	6/5 to 7/8	4,868	1.85	246.78	0.094	34,433	13.09	9,837,960	3,739
	7/8 to 8/5	4,200	1.81	196.56	0.084	30,769	13.24	8,791,160	3,782
	8/5 to 9/2	4,262	1.77	180.69	0.076	33,036	13.74	9,438,835	3,925
	9/2 to 9/30	4,390	1.74	157.62	0.062	26,158	10.38	7,473,715	2,965
	9/30 to 11/5	5,565	1.73	211.47	0.065	9,822	3.06	2,806,350	873
	Average		1.80		0.082		11.78		3,364
	Total	27,695	1	,255.96		174,587		49,882,120	

		Dry m (lb	atter s.)	Prot (1b	ein s.)	Caro (Mg	tene ms.)	Vitamin A eq (units	quivalen s)*
lot No.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
II	5/6 to 6/5	4,419	1.86	263.37	0.111	40,452	17.05	11,557,578	4,872
	6/5 to 7/8	4,884	1.85	247.62	0.094	34,549	13.06	9,871,290	3,732
	7/8 to 8/5	4,183	1.81	195.77	0.085	30,646	13.24	8,755,880	3,784
	8/5 to 9/2	4,242	1.78	179.86	0.075	32,884	13.82	9,395,424	3,948
	9/2 to 9/30	4,340	1.75	155.81	0.063	25,858	10.45	7,387,920	2,965
	9/30 to 11/5	5,495	1.74	208.81	0.067	9,699	3.07	2,771,050	878
	Average		1.80		0.082		11.78		3,363
	Total	27,563	l	,251.24		174,088		49,739,142	

		Dry m	atter s.)	Prot (1b	ein s.)	Caro (Mg	tene ms.)	Vitamin A ed	quivalent s)*
Lot No.	Dates	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day	Total	Av.per 100# per day
v	5/6 to 6/5	3,915	1.95	233.33	0.116	35,838	17.86	10,239,420	5,103
	6/5 to 7/8	4,366	1.87	221.35	0.096	30,884	13.26	8,824,167	3,788
	7/8 to 8/5	3,747	1.83	175.37	0.086	27,452	13.38	7,843,500	3,824
	8/5 to 9/2	3,805	1.79	161.34	0.076	29,498	13.86	8,427,974	3,960
	9/2 to 9/30	3,893	1.76	139.77	0.063	23,197	10.48	6,627,679	2,995
	9/30 to 11/5	4,930	1.75	187.33	0.066	8,701	3,08	2,486,003	880
	Average		1.82		0.084		11.99		3,425
	Total	24,656	l	,118.49		155,570		44,448,743	

Lot No.	Animal No.	12-12-46	2-5-47	4-2-47	5-14-47	6-12-47	8-7-47	10-2-47	11-13-47
I	2	90.90	87.00	131.40	90.00	588.00	513.00	462.00	168.00
	3	104.40		132.00	96.00	522.00	597.00	606.00	294.00
	7	30.00	111.00	108.00	82.50	591.60	597.00	588.00	183.00
	9	36.00		105.00	57.00	534.00	366.00	234.00	129.00
	10	50.40	\$7.00	87.00	67.50	500.40	644.50	372.30	174.00
	14	153.00	\$7.00	97.20	69.00	504.00	618.00	492.00	285.00
	17	50.10		123.00	\$3.10	420.60	498.00	575.40	210.00
	23	47.40	63.00	114.00	90.00	498.00	561.00	324.00	246.00
	30	57.00		66.00	60.00	431.40	552.00	315.00	150.00
	43	57.00		112.50	87.00	522.00	561.00	552.00	399.00
Avera	ge	67.62	87.00	107.61	78.21	511.20	550.75	452.07	223.80

Lot No.	Animal No.	12-12-46	2-5-47	4-2-47	5-14-47	6-12-47	8-7-47	10-2-47	11-13-47
II	5	50.70	105.00	114.00	57.00	393.00	543.00	618.00	447.00
	6	50.70	57.00	\$7.00	69.00	393.00	579.00	492.00	159.00
	18	24.00	45.00	129.00	102.00	431.40	477.00	294.00	189.00
	21	26.40	75.00	84.00	48.00	381.00	429.00	348.00	129.00
	24	93.00	89.40	114.00	92.40	584.40	597.00	411.00	225.00
	25	85.20	31.80	78.00	87.00	498.00	494.40	522.00	150.00
	33	37.50	63.00	69.00	67.50	524.40	561.00	300.00	135.00
	42	63.00	69.00	75.00	84.00	462.00	570.00	300.00	276.00
	39	63.00	55.20	81.00	67.50	518.40	504.00	267.00	171.00
	40	71.40	69.00	96.00	91.50	600.60	720.00	747.00	261.00
Averag	ge	56.49	65.94	92.70	76.59	478.62	547.44	429.90	214.20

÷.

Lot No.	Animal No.	12-12-46	2-5-47	4-2-47	5-14-47	6-12-47	8-7-47	10-2-47	11-13-47
III	1	88.20	135.00	175.80	132.00	534.00	522.00	330.00	222.00
	11	59.40		111.00	114.00	348.00	276.00	331.20	219.00
	12	45.00	108.00	135.00	102.00	348.00	462.00	330.00	183.00
	13	72.60	177.00	156.00	108.00	524.40	351.00	513.00	201.00
	15	60.00	192.00	184.20	162.00	618.00	688.50	315.00	234.00
	27	56.40	133.20	126.00	102.00	507.60	513.00	294.00	156.00
	29	67.80	103.80	132.00	136.20	469.50	469.50	597.00	210.00
	31	73.20	103.80	162.00	135.00	543.00	387.00	462.00	225.00
	36	88.20	156.00	123.00	105.00	552.00	552.00	606.00	522.00
	37	69.60	120.30	144.00	109.80	552.00	405.00	528.00	153.00
lvera	ge	68.04	136.57	144.90	120.60	499.65	462.60	430.62	232.50

Lot No.	Animal No.	12-12-46	2-5-47	4-2-47	5-14-47	6-12-47	8-7-47	10-2-47	11-13-47
IV	4	48.00	54.00	252.00	451.50	618.00	469.50	534.00	504.00
	8	65.40	67.20	277.50	484.50	579.00	582.60	345.00	243.00
	16	111.00		316.20	513.00	588.00	552.00	561.00	246.00
	19	57.00	41.10	316.20	423.00	575.40	504.00	351.00	238.50
	20	51.00	70.80	228.00	405.00	466.50	351.00	447.00	
	22	48.90	81.00	339.00	570.00	702.00	717.00	675.00	267.00
	26	48.90		268.50	399.00	462.00	454.50	306.00	144.00
	28	154.80	124.80	318.00	570.00	762.00	747.00	606.00	333.00
	32	52.20	66.00	246.00	450.00	435.00	447.00	264.00	174.00
	34	49.80	66.00	264.00	528.00	610.80	570.00	579.00	174.00
Avera	ge	68.70	71.36	282.54	479.40	579.87	539.46	466.80	258.16

X

Lot No.	Animal No.	11-13-47	1-8-48	3-4-48	5-6-48	7-8-48	9-2-48	11-5-48
I	2	168.00	129.00	117.00	40.50	496.50	441.00	63.00
	3	294.00	198.00	135.00	47.40	876.00	571.50	220.50
	7	183.00	160.20	72.00	43.50	585.00	475.50	63.00
	9	129.00	138.00	66.00	49.50	369.00	294.00	51.00
	10	174.00	138.00	76.50	38.40	597.00	391.50	66.00
	14	285.00	96.00	81.00	43.50	609.00	648.00	97.50
	17	210.00	191.70	102.00	38.40	648.00	609.00	61.50
	23	246.00	121.20	96.00	36.90	723.00	496.50	76.50
	30	150.00	57.00	82.50	30.60	747.00	369.00	57.00
	43	399.00	159.00	102.00	28.50	723.00	648.00	150.00
Averag	е	223.80	138.81	93.00	39.72	637.35	494.40	90.60

Lot No.	Animal No.	11-13-47	1-8-48	3-4-48	5-6-48	7-8-48	9-2-48	11-5-48
II	5	447.00	82.50	90.00	31.80	507.00	519.00	159.00
	6	159.00	67.50	91.50	28.80	648.00	432.00	76.50
	18	189.00	93.00	156.00	34.20	465.00	330.00	60.00
	21	129.00	109.80	96.00	28.50	414.00	324.00	94.50
	24	225.00	145.20	96.00	36.90	723.00	681.00	138.00
	25	150.00	82.50	112.50	33.00	475.50	399.00	61.50
	33	135.00	76.50	82.50	29.40	519.00	406.50	117.00
	42	276.00	123.00	111.00	66.00	571.50	486.00	78.00
	39	171.00	92.40	112.50	31.80	507.00	441.00	117.00
	40	261.00	84.00	90.00	38.40	939.00	747.00	90.00
lvera	ge	214.20	95.64	103.80	35.88	576.90	476.55	99.15

Lot No.	Animal No.	11-13-47	1-8-48	3-4-48	5-6-48	7-8-48	9-2-48	11-5-48
III	1	222.00	141.00	171.00	49.50	558.00	399.00	60.00
	11	219.00	102.00	144.00	31.80	349.50	300.00.	45.00
	12	183.00	117.00	144.00	42.00	519.00	609.00	58.50
	13	201.00	120.00	150.00	38.40	609.00	450.00	57.00
	15	234.00	117.00	150.00	35.40	771.00	1080.00	1125.00
	27	156.00	108.00	147.00	43.50	747.00	771.00	76.50
	29	210.00	174.00	135.00	43.50	648.00	531.00	60.00
8	31	225.00	138.00	123.00	36.90	609.00	585.00	58.50
	36	522.00	165.00	120.00	46.50	648.00	539.10	148.50
	37	153.00	150.00	144.00	46.50	648.00	539.10	72.00
Averag	е	232.50	133.20	142.80	41.40	610.65	580.32	176.10

Lot No.	Animal No.	11-13-47	1-8-48	3-4-48	5-6-48	7-8-48	9-2-48	11-5-48
IV	4	504.00	168.00	46.20	492.90	939.00	747.00	66.00
	8	243.00	255.00	76.50	369.00	967.50	681.00	55.50
	16	246.00	207.00	67.50	492.90	822.00	571.50	115.20
	19	238.50	261.00	93.00	321.90	681.00	519.00	60.00
	22	267.00	417.00	135.00	458.70	1110.00	1110.00	159.00
	26	144.00	135.00	51.00	426.00	597.00	496.50	84.00
	28	333.00	405.00	147.00	495.00	996.00	1023.00	70.50
	32	174.00	225.00	66.00	369.00	681.00	747.00	61.50
	34	174.00	280.50	93.00	513.60	1080.00	558.00	75.00
Avera	ge	258.16	261.50	86.13	437.66	874.83	717.00	82.97

Lot No.	Animal No.	12-12-46	2-5-47	4-2-47	5-14-47	6-12-47	8-7-47	10-2-47	11-13-47
I	2		185.40	114.00	171.00	323.40	255.00	291.60	368.40
	3			100.80	147.30	530.10	328.20	285.30	443.70
	7		153.00	119.70	159.60	438.90	289.20	311.40	322.50
	9				118.20	454.20	156.30	286.20	354.60
	10		173.10	100.80	160.20	539.70	300.90	276.60	353.70
	14		168.90	93.30	148.80	520.20	299.40	249.60	342.60
	17			123.00	160.80	230.70	275.40	317.10	318.00
	23		200.10	89.40	168.00	429.30	260.40	311.70	465.30
	30			116.70	127.50	393.90	288.60	2 63.40	418.50
	43			134.10	169.20	395.10	218.40	288.60	357.30
Avera	ge		176.10	100.87	153.06	425.55	267.18	288.15	374.46

Lot No.	Animal No.	12-12-46	2-5-47	4-2-47	5-14-47	6-12-47	8-7-47	10-2-47	11-13-47
II	5		115.50	105.30	139.20	322.50	241.50	378.90	387.00
	6		214.80	131.40	169.80	280.50	218.10	435.60	495.30
	18		222.00	101.70	222.60	330.90	316.20	404.70	491.10
	21		147.30		179.40	471.90	535.80	380.40	474.00
	24	128.40	143.70	131.70	138.30	336.90	230.70	372.90	423.90
	25	108.90	165.30	120.00	162.00	424.80	221.40	341.10	364.50
	33	141.60	191.40	93.90	113.70	468.90	191.40	351.00	260.10
	42	193.50	151.50	141.00	134.70	305.10	229.50	336.00	333.30
	39	186.30	201.90	74.70	142.20	366.60	217.20	537.00	401.40
	40	126.60	224.10	142.20	174.90	458.70	276.00	479.70	242.70
Aver	age	147.55	177.75	115.77	157.68	376.68	272.28	401.73	387.33

Lot No.	Animal No.	12-12-46	2-5-47	4-2-47	5-14-47	6-12-47	8-7-47	10-2-47	11-13-47
III	1		150.00	152.70	189.60	464.70	222.60	307.50	437.10
	11			105.90	103.20	254.40	295.80	321.90	426.30
	12	176.10	161.10	94.20	134.10	527.40	212.10	426.00	322.50
	13	238.50	158.70	121.50	167.40	633.90	142.20	426.00	552.90
	15	165.00	198.30	77.70	117.60	648.30	333.60	416.10	524.70
	27	119.40	162.00	86.40	126.60	386.70	174.00	443.70	415.80
	29	163.20	162.00	145.50	129.30	445.80	201.30	268.20	274.50
	31	234.00	207.90	186.00	207.90	919.80	217.50	350.10	435.60
	36	225.90	198.30	106.80	153.60	422.10	271.80	303.30	482.10
	37	133.80	182.70	111.00	166.50	567.60	365.10	392.40	417.00
ver	age	181.99	175.67	118.77	149.58	527.07	243.60	365.52	428.85

Lot No.	Animal No.	12-12-46	2-5-47	4-2-47	5-14-47	6-12-47	8-7-47	10-2-47	11-13-47
εv	4		149.40	154.20	354.90	468.90	208.80	416.70	457.20
	8		141.90	174.90	369.90	461.40	181.80	536.70	390.00
	16			126.00	357.00	365.40	195.60	431.40	402.30
	19	233.40	191.40	132.00	295.80	317.10	230.70	499.20	417.60
	20	171.60	126.60	104.40	288.60	289.50	320.40	483.00	Died
	22	182.40	131.40	166.80	373.50	486.60	292.20	566.10	324.00
	26	175.20		183.60	291.60	557.10	229.80	492.30	340.20
	28	202.80	188.60	185.70	448.50	891.90	321.00	342.30	438.00
	32	\$1.60	127.80	155.70	310.50	392.40	204.00	418.20	434.70
	34	174.90	212.70	184.80	350.40	442.20	343.50	444.60	392.70
ver	age	174.55	158.70	156.81	344.07	467.25	252.78	463.05	399.63

Lot No.	Animal No.	11-13-47	1-8-48	3-4-48	5-6-48	7-8-48	9-2-48	11-5-48
I	2	368.40	454.80	295.20	242.70	433.50	485.40	519.30
	3	443.70	435.90	200.10	194.70	505.80	372.90	458.70
	7	322.50	414.00	236.10	189.00	380.10	299.10	457.80
	9	354.60	442.80	402.30	226.80	370.80	245.70	490.20
	10	353.70	462.90	444.00	209.10	533.70	381.90	483.30
	14	342.60	372.30	449.40	229.50	454.80	392.40	388.20
	17	318.00	413.10	345.60	219.60	392.40	292.80	443.40
	23	465.30	416.40	514.80	259.50	490.50	364.50	385.50
	30	418.50	310.20	283.80	186.00	480.00	202.80	386.70
	43	357.30	414.30	290.10	188.10	304.50	275.40	376.50
lverag	ge	374.46	413.67	346.14	214.50	434.61	331.29	438.96

Lot No.	Animal No.	11-13-47	1-8-48	3-4-48	5-6-48	7-8-48	9-2-48	11-5-48
II	5	387.00	420.30	507.00	321.60	477.00	396.60	399.60
	6	495.30	361.20	287.40	224.10	425.40	318.60	325.50
	18	491.10	444.00	373.80	260.70	447.60	330.00	309.00
	21	474.00	475.50	361.80	224.10	389.70	259.20	389.40
	24	423.90	452.40	292.80	232.50	439.50	218.40	450.90
	25	364.50	354.30	327.30	234.00	416.10	333.60	471.90
	33	260.10	378.00	250.80	213.30	513.60	229.50	391.20
	42	333.30	423.00	249.90	306.30	372.90	280.80	384.90
	39	401.40	436.80	507.30	201.60	243.00	213.90	418.20
	40	242.70	440.70	307.50	270.60	324.60	321.00	445.50
Avera	ge	387.33	418.62	346.56	248.88	404.94	290.16	398.61

Lot No.	Animal No.	11-13-47	1-8-48	3-4-48	5-6-48	7-8-48	9-2-48	11-5-48
III	1	437.10	449.40	291.90	313.80	537.90	320.10	301.50
	11	426.30	425.10	413.70	246.60	321.30	270.00	357.60
	12	322.50	446.70	379.20	341.10	450.60	273.30	417.60
	13	552.90	417.00	432.00	282.60	454.80	276.00	379.20
	15	524.70	425.70	445.50	221.10	508.20	330.00	440.10
	27	415.80	329.40	338.70	217.50	294.00	241.20	325.50
	29	274.50	421.20	249.60	229.50	392.40	319.20	510.00
	31	435.60	395.40	265.50	298.20	541.80	152.10	405.60
	36	482.10	450.60	282.00	279.00	455.40	414.30	446.10
	37	417.00	469.50	493.20	279.00	402.90	333.30	525.60
Averag	ge	428.85	423.00	359.13	270.84	435.93	292.95	410.88

Lot No.	Animal No.	11-13-47	1-8-48	3-4-48	5-6-48	7-8-48	9-2-48	11-5-48
IV	4	457.20	471.90	205.80	663.30	494.10	163.50	427.80
	8	390.00	432.60	241.50	477.30	293.70	179.40	512.10
	16	402.30	385.20	269.70	678.30	419.10	330.90	353.10
	19	417.60	395.40	268.50	455.10	387.90	354.60	309.00
	22	324.00	325.20	371.10	521.40	460.50	400.50	386.10
	26	340.20	267.60	252.90	451.80	425.70	324.00	322.20
	28	438.00	317.10	339.00	755.10	559.80	210.30	337.20
	32	434.70	398.10	207.30	421.80	387.90	189.00	282.90
	34	392.70	421.20	234.00	789.00	354.00	261.90	371.10
Avera	ge	399.63	379.37	265.53	579.23	420.30	268.23	366.83

Lot No.	Animal No.	12-12	2-5	4-2	5-14	6-12	8-7	10-2	11-13
I	2	7.94	8.75	9.80	11.90	7.34	6.96	12.30	10.90
	3	5.30	5.81	6.60	9.95	8.31	5.60	6.38	8.06
	7	6.68	6.40	8.32	11.60	8.00	7.70	7.25	8.80
	9	5.40	5.15	8.40	10.25	8.44	9.35	7.60	8.25
	10	5.10	7.40	5.80	8.44	6.60	7.30	5.63	6.75
	14	7.70	6.56	6.60	8.86	5.30	8.25	7.38	9.05
	17	5.30	4.50	6.60	9.65	8.35	10.25	8.04	8.70
	23	5.10	6.50	6.80	10.25	7.85	9.11	7.68	10.25
	30	8.54	5.55	9.20	9.05	6.75	9.50	9.32	8.35
	43	5.96	8.40	8.50	10.10	7.55	10.75	8.60	8.00
Averag	e	6.30	6.50	7.66	10.00	7.45	8.48	8.02	8.71

Lot No.	Animal No.	12-12	2-5	4-2	5-14	6-12	8-7	10-2	11-13
II	5	6.40	7.59	10.00	9.65	5.70	10.75	4.13	7.40
	6	6.40	7.65	10.40	10.40	6.75	8.80	5.38	9.65
	18	5.40	8.19	9.92	10.25	6.90	9.35	8.45	9.65
	21	7.10	7.05	7.00	10,10	8.35	7.64	6.10	7.00
	24	6.90	5.95	8.40	10.60	6.60	7.10	7.38	8.50
	25	5.10	3.81	8.70	10.19	8.90	8.90	8.60	8.35
	33	5.40	5.45	8.50	10.25	7.40	8.00	5.80	8.90
	42	6.52	7.98	9.00	10.75	7.30	8.35	7.15	7.30
	39	6.48	5.45	10.20	10.34	9.35	8.90	6.95	7.14
	40	8.20	7.30	9.38	11.25	8.15	9.59	6.68	8.15
Averag	e	6.39	6.64	9.15	10.38	7.54	8.74	6.66	8.20

Lot No.	Animal No.	12-12	2-5	4-2	5-14	6-12	8-7	10-2	11-13
III	1	6.60	6.40	8.70	8.25	7.70	7.55	8.00	9.35
	11	8.20	6.93	7.20	9.65	7.20	7.49	8.00	8.25
	12	8.40	6.05	8.40	9.35	6.45	10.40	8.00	8.70
	13	6.66	3.75	8.70	8.80	7.00	10.60	4.53	6.45
	15	7.10	2.95	8.00	9.83	7.30	7.85	7.25	6.75
	27	5.90	5.48	9.20	10.34	6.75	9.65	7.80	7.85
	29	8.32	9.00	9.20	10.34	9.35	8.50	6.85	8.35
	31	6.70	7.65	9.20	10.60	7.10	11.40	8.60	9.35
	36	6.82	6.85	7.80	9.35	6.75	11.10	8.45	6.75
	37	6.70	8.00	7.80	10.10	8.70	8.90	6.68	8.00
Averag	ge	7.14	6.31	8.42	9.66	7.43	9.34	7.42	7.98

Lot No.	Animal No.	12-12	2-5	4-2	5-14	6-12	8-7	10-2	11-13
IV	4	7.54	6.15	7.20	8.50	5.80	8.90	7.68	8.00
	8	6.20	3.65	7.60	9.05	9.80	8.80	7.25	7.55
	16	5.30	5.25	6.60	9.95	9.20	7.24	5.95	7.40
	19	6.60	4.40	8.40	9.95	10.10	8.70	7.08	7.30
	20	6.70	7.05	8.70	9.05	8.25	8.70	9.10	
	22	5.52	5.29	6.30	9.65	10.75	9.61	7.25	8.25
	26	6.66	5.35	6.50	8.50	7.70	9.20	7.90	7.20
	28	6.66	7.30	8.50	10.40	8.80	7.20	8.60	9.53
	32	7.10	7.30	7.80	8.70	6.75	8.25	6.85	7.10
	34	8.20	4.60	7.40	8.70	7.30	6.45	5.63	7.00
Averag	ge	6.65	5.63	7.50	9.24	8.44	8.30	7.33	7.70

Lot No.	Animal No.	11-13	1-8	3-4	5-6	7-8	9-2	11-5
I	2	10.90	11.25	8.25	4.70	7.05	7.35	4.50
	3	8.06	14.85	6.35	5.25	7.15	6.90	4.50
	7	8.80	8.35	7.00	6.50	7.90	7.35	4.90
	9	8.25	9.35	8.25	6.80	6.65	7.25	3.20
	10	6.75	7.06	6.35	5.90	6.80	6.90	2.75
	14	9.05	9.35	6.25	6.80	7.60	6.65	6.25
	17	8.70	8.90	7.40	6.96	8.60	8.00	5.25
	23	10.25	8.21	6.75	6.11	7.60	7.05	4.15
	30	8.35	10.90	7.85	6.11	5.80	7.45	5.70
	43	8.00	10.10	7.55	6.65	7.60	8.00	4.80
Averag	e	8.71	9.83	7.20	6.18	7.28	7.29	4.60

Lot No.	Animal No.	11-13	1-8	3-4	5-6	7-8	9-2	11-5
II	5	7.40	9.35	6.60	6.96	5.25	6.59	4.80
	6	9.65	7.16	6.75	7.05	7.25	5.90	4.35
	18	9.65	8.76	8.25	6.99	7.90	6.90	5.00
	21	7.00	8.74	6.60	7.11	6.96	7.05	3.45
	24	8.50	8.25	6.60	4.05	10.85	7.60	4.15
	25	8.35	8.80	7.10	4.80	6.05	7.35	4.15
	33	8.90	9.35	6.05	7.15	7.90	7.90	3.65
9	42	7.30	10.40	5.70	7.35	9.05	6.65	5.25
	39	7.14	8.80	6.25	6.65	6.90	6.80	4.80
	40	8.15	12.05	8.00	7.39	7.15	8.00	5.15
lverage	)	8.20	9.17	6.79	6.55	7.53	7.07	4.48

Lot No.	Animal No.	11-13	1-8	3-4	5-6	7-8	9-2	11-5
III	1	9.35	9.65	5.99	7.05	4.80	6.05	5.00
	11	8.25	7.70	8.00	7.45	7.75	7.35	7.05
	12	8.70	7.06	7.20	6.80	6.90	6.90	5.15
	13	6.45	8.00	6.35	6.05	7.25	6.05	3.95
	15	6.75	6.60	5.00	6.65	6.05	7.25	3.10
	27	7.85	6.07	5.30	6.80	7.25	7.25	4.90
	29	8.35	9.80	7.40	7.60	7.15	7.60	6.15
	31	9.35	9.80	6.66	7.31	6.90	6.65	5.00
	36	6.75	5.80	7.40	7.31	6.05	5.60	3.75
	37	8.00	9.80	7.40	7.31	8.15	6.90	5.70
Averag	ge	7.98	8.03	6.67	7.03	6.83	6.76	4.98

Lot No.	Animal No.	11-13	1-8	3-4	5-6	7-8	9-2	11-5
IV	4	8.00	8.00	7.10	7.90	6.90	8.00	5.00
	8	7.55	8.00	7.10	7.90	6.90	5.90	4.60
	16	7.40	8.80	7.10	8.03	6.15	7.25	5.80
	19	7.30	8.80	6.45	7.75	4.80	7.15	4.86
	22	8.25	9.80	7.85	8.15	6.59	7.25	3.95
	26	7.20	11.66	8.00	8.15	7.25	8.25	5.25
	28	9.53	8.25	8.44	8.40	6.40	6.40	5.00
	32	7.10	7.70	5.30	7.75	5.50	5.60	3.85
	34	7.00	8.00	5.30	7.31	4.80	6.05	3.65
Averag	çe	7.70	8.78	6.26	7.93	6.14	6.87	4.66

Lot No.	Animal No.	12-12	2-5	4-2	5-14	6-12	8-7	10-2	11-13
I	2	7.65	7.55	7.60	7.05	7.35	6.20	8.55	9.35
	3		7.40	7.85	8.40	10.05	7.45	11.80	9.20
	7	7.20		7.90	6.25	9.35	7.20	7.70	8.15
	9	8.30	7.20	12.40	7.75	7.20	6.45	8.10	8.85
	10	8.45		7.45	7.45	11.30	7.25	4.20	9.65
	14	8.45	7.25	8.35	8.05	7.40	7.30	9.20	9.70
	17	7.70	7.60	8.10	7.25	11.25	6.65	8.35	10.30
	23	7.40	7.55	9.65	6.90	8.95	7.05	7.15	10.10
	30	7.95	8.00	10.35	6.75	10.55	7.65	13.45	10.55
	43	7.75	7.48	8.15	7.15	10.25	7.35	11.70	7.25
Avera	ge	7.87	7.50	8.78	7.30	9.36	7.06	9.02	9.31

Lot No.	Animal No.	12-12	2-5	4-2	5-14	6-12	8-7	10-2	11-13
II	5	6.05	7.20	13.15	9.20	10.20	7.75	9.60	9.00
	6	7.25		8.30	6.95	9.15	7.30	13.10	8.60
	18	7.05	7.50	8.20	7.05	8.65	7.45	14.00	9.00
	21	7.95		10.10	8.80	10.05	7.20	14.40	8.15
	24	6.85	7.85	8.70	6.85	8.50	7.30	15.80	8.45
	25	8.30	7.55	8.75	8.15	8.95	8.10	7.85	9.35
	33	8.30	7.20	8.30	8.40	7.85	7.95	9.65	7.70
	42	7.25	7.20	9.85	7.65	8.80	8.35	8.35	8.90
	39	7.75	7.25	8.35	6.95	11.25	6.55	10.10	10.55
	40	13.10	6.40	9.50	7.25	9.15	7.20	15.05	7.90
Averag	;e	7.98	7.27	9.32	7.72	9.26	7.52	11.79	8.76

Lot No.	Animal No.	12-12	2-5	4-2	5-14	6-12	8-7	10-2	11-13
III	l	6.60	8,00	8.60	8.40	9.15	6.65	13.05	9.55
	11	7.50	7.25	8.60	7.25	11.20	7.95	6.80	8.60
	12	7.60	7.00	8.20	7.65	7.90	6.30	9.65	9.70
	13	8.30	8.10	7.85	8.20	9.65	7.65	11.15	9.35
	15	8.10		11.70	8.40	9.25	8.35	7.60	9.70
	27	8.25	7.70	8.00	7.65	11.25	7.95	8.20	9.40
	29	7.75		8.05	7.25	8.45	8.35	8.55	8.65
	31	7.75	6.10	8.90	8.05	8.45	6.95	8.10	8.20
	36	6.90		9.05	7.20	7.75	7.95	7.60	8.85
	37	7.60	7.25	8.05	8.55	10.40	8.20	7.90	8.40
Averag	е	7.64	7.34	8.70	7.86	9.34	7.63	8.86	9.04
Lot No.	Animal No.	12-12	2-5	4-2	5-14	6-12	8-7	10-2	11-13
------------	---------------	-------	------	-------	------	-------	-------	------	-------
IV	4	8.30	7.20	7.30	9.25	8.50	8.55	8.25	9.10
	8	8.45	8.60	10.35	8.85	7.85	8.30	9.40	9.05
	16	7.95	8.25	8.75	9.15	10.85	8.10	9.00	9.00
	19	7.40	8.00	7.75	8.25	9.80	7.95	7.15	9.00
	20	8.95		8.30	8.70	9.35	8.20	8.15	
	22	8.25		8.20	8.55	9.15	7.95	7.05	9.35
	26	8.85	8.00	8.20	9.00	9.15	8.00	7.40	9.00
	28	7.40	7.00	7.40	8.25	10.25	7.40	7.15	8.30
	32	9.15		10.30	8.55	9.75	8.25	6.00	8.90
	34	10.00	7.95	9.50	7.65	8.15	7.95	7.40	8.75
Avera	ge	8.47	7.86	8.60	8.62	9.28	\$.06	7.70	\$.94

Lot No.	Animal No.	11-13	1-8	3-4	5-6	7-8	9-2	11-5
I	2	9.35	8.30	8.50	8.80	8.85	8.10	12.60
	3	9.20	9.40	8.25	9.60	8.95	10.10	13.10
	7	8.15	8.60	8.70	8.85	7.85	8.40	12.45
	9	8.85	8.35	8.50	9.40	8.90	8.25	13.00
)	10	9.65	9.35	8.75	9.20	8.00	9.25	13.00
,	14	9.70	8.85	8.65	.9.10	8.65	9.60	13.10
(	17	10.30	9.20	8.50	9.65	8.50	9.50	13.30
	23	10.10	9.00	8.05	8.70	9.80	7.95	12.45
	30	10.55	8.75	7.75	8.90	8.60	9.65	13.00
	43	7.25	9.20	8.55	8.60	7.85	8.40	12.30
Ave	rage	9.31	8.90	8.42	9.08	8.60	8.92	12.83

Lot No.	Animal No.	11-13	1-8	3-4	5-6	7-8	9-2	11-5
II	5	9.00	9.20	9.70	9.35	8.85	9.40	12.30
	6	8.60	9.20	8.80	8.95	8.90	8.25	11.40
	18	9.00	8.35	8.65	9.00	8.30	8.50	12.15
	21	8.15	10.00	9.05	9.00	7.85	8.15	12.30
	24	8.45	9.00	9.90	9.00	8.25	8.10	12.20
	25	9.35	8.55	9.65	9.05	8.10	9.35	12.15
	33	7.70	9.65	8.65	8.80	8.85	8.35	12.95
	42	8.90	9.30	9.45	10.05	8.70	9.00	13.40
	39	10.55	9.20	10.95	9.00	9.75	10.00	13.30
	40	7.90	8.35	10.00	9.45	8.35	8.60	12.55
Averag	ge	8.76	9.08	9.48	9.16	8.59	8.77	12.47

Lot No.	Animal No.	11-13	1-8	3-4	5-6	7-8	9-2	11-5
III	l	9.55	9.25	9.20	9.05	8.65	9.05	14.35
	11	8.60	8.65	9.70	9.45	7.75	9.40	12.30
	12	9.70	9.00	9.75	11.05	9.60	9.00	12.50
	13	9.35	9.20	9.95	10.05	8.75	9.05	13.60
	15	9.70	9.15	9.20	9.60	9.10	9.10	13.85
	27	9.40	9.20	9.20	9.70	9.20	8.90	12.65
	29	8.65	9.20	8.80	11.40	7.70	10.15	13.05
	31	8.20	9.50	8.65	10.10	9.30	9.95	11.90
	36	8.85	8.55	8.20	10.20	8.65	8.55	12.00
	37	8.40	8.95	8.65	9.90	9.10	9.10	12.25
Averag	ge	9.04	9.06	9.13	10.05	8.78	9.22	12.84

Lot No.	Animal No.	11-13	1-8	3-4	5-6	7-8	9-2	11-5
IV	4	9.10	9.25	8.60	10.40	8.45	9.50	13.95
	8	9.05	9.00	8.40	11.15	8.90	9.95	12.25
	16	9.00	9.20	9.20	10.20	8.20	9.95	12.45
	19	9.00	9.00	8.50	10.65	8.75	9.25	12.85
	22	9.35	9.10	9.20	11.30	8.50	9.70	12.90
	26	9.00	8.80	9.65	10.60	8.50	9.25	12.35
	28	8.30	9.15	8.90	8.55	8.80	9.65	12.10
	32	8.90	8.50	8.70	10.90	8.45	10.15	12.45
	34	8.75	8.90	9.00	10.95	8.25	9.05	13.90
Avera	ge	8.94	8.99	8.91	10.52	8.53	9.61	12.80

lot	Animal No.	12-11-12 1946	1-8-47	2-5-47	3-5-47	4-2-47	4-30-47	5-14-15 1947	Total gain
I	2	460.00	490.00	545.00	565.00	600.00	605.00	590.00	130.00
	3	342.50	375.00	425.00	445.00	470.00	505.00	510.00	167.50
	7	267.50	305.00	340.00	345.00	380.00	395.00	397.50	130.00
	9	232.50	250.00	295.00	320.00	355.00	380.00	377.50	145.00
	10	327.50	260.00	295.00	315.00	320.00	335.00	330.00	92.50
	14	285.00	305.00	330.00	315.00	340.00	325.00	317.50	32.50
	17	285.00	290.00	345.00	365.00	390.00	410.00	410.00	125.00
	23	385.00	420.00	470.00	495.00	525.00	545.00	550.00	165.00
	30	290.00	295.00	325.00	335.00	355.00	355.00	325.00	35.00
	43	250.00	280.00	335.00	375.00	410.00	435.00	425.00	175.00
Avei	rage	303.50	327.00	370.50	387.50	414.50	429.00	423.25	119.75

Lot No.	Animal No.	12-11-12 1946	1-8-47	2-5-47	3-5-47	4-2-47	4-30-47	3-14-15 1947	Total gain
II	5	320.00	340.00	390.00	405.00	450.00	475.00	442.50	122.50
	6	280.00	315.00	360.00	375.00	405.00	420.00	427.50	147.50
	18	355.00	400.00	460.00	475.00	500.00	505.00	522.50	167.50
	21	332.50	365.00	420.00	440.00	475.00	480.00	482.50	150.00
	24	282.50	295.00	330.00	360.00	375.00	395.00	390.00	107.50
	25	395.00	420.00	475.00	495.00	530.00	545.00	537.50	142.50
	33	265.00	290.00	320.00	355.00	370.00	400.00	395.00	130.00
	42	240.00	250.00	290.00	300.00	330.00	335.00	342.50	102.50
	39	367.50	380.00	420.00	440.00	465.00	495.00	482.50	115.00
	40	215.00	235.00	265.00	280.00	305.00	330.00	307.50	92.50
lvera	lge	305.25	329.00	373.00	392.50	420.50	438.00	433.00	127.75

Lot No.	Animal No.	12-11-12 1946	1-8-47	2-5-47	3-5-47	4-2-47	4-30-47	5-14-15 1947	Total gain
III	1	350.00	385.00	430.00	450.00	470.00	500.00	495.00	145.00
	11	285.00	305.00	345.00	370.00	400.00	410.00	412.50	127.50
	12	235.00	260.00	295.00	315.00	345.00	365.00	360.00	125.00
	13	422.50	470.00	530.00	555.00	580.00	590.00	582.50	160.00
	15	262.50	280.00	315.00	330.00	360.00	395.00	385.00	122.50
	27	245.00	265.00	300.00	330.00	350.00	375.00	362.50	117.50
	29	295.00	325.00	370.00	405.00	440.00	470.00	472.50	177.50
	31	317.50	345.00	385.00	400.00	435.00	445.00	445.00	127.50
	36	347.50	375.00	410.00	425.00	440.00	460.00	447.50	100.00
	37	282.50	310.00	330.00	350.00	375.00	395.00	392.50	110.00
Average	e	304.25	332.00	371.00	393.00	419.50	440.50	435.50	131.25

Lot No.	Animal No.	12-11-12 1946	1-8-47	2-5-47	3-5-47	4-2-47	4-30-47	5-14-15 1947	Total gain
IV	4	315.00	325.00	340.00	355.00	365.00	400.00	400.00	\$5.00
	8	257.50	280.00	295.00	315.00	335.00	365.00	375.00	117.50
	16	395.00	410.00	425.00	445.00	470.00	510.00	522.50	127.50
	19	247.50	280.00	285.00	310.00	325.00	360.00	365.00	117.50
	20	300.00	305.00	310.00	345.00	350.00	390.00	400.00	100.00
	22	280.00	290.00	295.00	315.00	350.00	385.00	385.00	105.00
	26	230.00	260.00	270.00	285.00	315.00	350.00	355.00	125.00
	28	272.50	280.00	275.00	285.00	300.00	315.00	327.50	55.00
	32	347.50	380.00	375.00	400.00	420.00	445.00	452.50	105.00
	34	282.50	310.00	325.00	345.00	375.00	390.00	392.50	110.00
Averag	e	292.75	312.00	319.50	340.00	360.50	391.00	397.50	104.75

 $\overline{\alpha}$ 

Lot No.	Animal No.	5-14-15 1947	6-12-47	7-10-47	8-7-47	9-4-47	10-2-47	10-30-47	11-13-14 1947	Total gain
I	2	590.00	638.00	664.00	720.00	786.00	740.00	747.00	734.50	144.50
	3	510.00	520.00	540.00	609.00	654.00	646.00	655.00	655.00	145.00
	7	397.50	416.00	456.00	528.00	582.00	565.00	590.00	565.00	167.50
	9	377.50	410.00	428.00	494.00	540.00	580.00	562.00	570.00	192.50
	10	330.00	346.00	367.00	425.00	454.00	478.00	476.00	442.00	112.00
	14	317.50	314.00	354.00	382.00	394.00	388.00	400.00	388.00	70.50
	17	410.00	424.00	480.00	538.00	564.00	587.00	576.00	568.50	158.50
	23	550.00	544.00	604.00	670.00	714.00	760.00	743.00	731.00	181.00
	30	325.00	320.00	378.00	430.00	444.00	472.00	440.00	442.50	117.50
	43	425.00	440.00	482.00	549.00	580.00	600.00	590.00	614.00	189.00
Avei	rage	423.25	437.20	475.30	534.50	571.20	581.60	577.90	571.05	147.80

Lot No.	Animal No.	5-14-15 1947	6-12-47	7-10-47	8-7-47	9-4-47	10-2-47	10-30-47	11-13-14 1947	Total gain
II	5	442.50	472.00	508.00	581.00	600.00	604.00	585.00	614.00	171.5
	6	427.50	422.00	472.00	524.00	526.00	574.00	582.00	557.50	130.0
	18	522.50	520.00	568.00	636.00	674.00	708.00	704.00	677.00	154.5
	21	482.50	506.00	538.00	612.00	654.00	645.00	688.00	660.00	177.5
	24	390.00	394.00	434.00	503.00	530.00	565.00	577.00	553.00	163.0
	25	537.50	548.00	580.00	644.00	674.00	670.00	684.00	690.00	152.5
	33	395.00	432.00	464.00	512.00	564.00	590.00	588.00	581.50	186.5
	42	342.50	342.00	406.00	460.00	492.00	524.00	507.00	495.00	152.5
	39	482.50	508.00	527.00	595.00	624.00	625.00	658.00	648.00	165.5
	40	307.50	308.00	353.00	410.00	440.00	424.00	440.00	418.50	111.0
Ave	rage	433.00	445.20	485.00	547.70	577.80	592.90	601.30	589.45	156.4

Lot No.	Animal No.	5-14-15 1947	6-12-47	7-10-47	8-7-47	9-4-47	10-2-47	10-30-47	11-13-14 1947	Total gain
III	1	495.00	510.00	556.00	606.00	640.00	702.00	714.00	699.00	204.00
	11	412.50	416.00	447.00	499.00	530.00	540.00	564.00	556.00	143.50
	12	360.00	382.00	440.00	485.00	534.00	534.00	570.00	537.00	177.00
	13	582.50	612.00	628.00	698.00	760.00	706.00	736.00	727.50	145.00
	15	385.00	400.00	427.00	484.00	510.00	542.00	520.00	506.00	121.00
	27	362.50	376.00	400.00	467.00	510.00	540.00	540.00	540.00	177.50
	29	472.50	486.00	508.00	560.00	600.00	594.00	620.00	597.00	124.50
	31	445.00	452.00	495.00	547.00	544.00	576.00	582.00	582.50	137.50
	36	447.50	478.00	534.00	598.00	608.00	622.00	621.00	640.00	192.50
	37	392.50	440.00	462.00	520.00	554.00	580.00	576.00	570.00	177.50
Avei	rage	435.50	455.20	489.70	546.40	579.00	593.60	604.30	595.50	160.00

Lot No.	Animal No.	5-14-15 1947	6-12-47	7-10-47	8-7-47	9-4-47	10-2-47	10-30-47	11-13-14 1947	Total gain
IV	4	400.00	446.00	482.00	547.00	560.00	584.00	591.00	590.00	190.00
	8	375.00	424.00	452.00	510.00	534.00	595.00	586.00	565.00	190.00
	16	522.50	580.00	610.00	710.00	744.00	742.00	780.00	747.50	225.00
	19	365.00	424.00	458.00	522.00	556.00	606.00	584.00	560.00	195.00
	20	400.00	428.00	493.00	554.00	580.00	595.00	596.00	Died	
	22	385.00	462.00	498.00	570.00	620.00	612.00	644.00	635.00	250.00
	26	355.00	396.00	440.00	496.00	526.00	560.00	540.00	536.00	181.00
	28	327.50	336.00	386.00	422.00	420.00	424.00	430.00	414.00	86.50
	32	452.50	480.00	544.00	611.00	656.00	690.00	640.00	637.00	184.50
	34	392.50	436.00	480.00	560.00	582.00	588.00	607.00	606.50	214.00
lvei	rage	397.50	441.20	484.30	550.20	577.80	599.60	599.80	587.88	190.66

Lot No.	Animal No.	11-13-14 1947	12-11-47	1-8-48	2-5-48	3-4-48	4-8-48	5-5-6 1948	Total gain
I	2	734.50	765.00	805.00	\$90.00	940.00	945.00	970.00	235.50
	3	655.00	700.00	715.00	755.00	780.00	815.00	\$65.00	210.00
	7	565.00	605.00	620.00	660.00	670.00	685.00	730.00	165.00
	9	570.00	610.00	630.00	695.00	725.00	755.00	780.00	210.00
	10	442.00	490.00	525.00	560.00	585.00	610.00	627.50	185.50
	14	388.00	440.00	470.00	500.00	525.00	555.00	575.00	187.00
	17	568.50	600.00	615.00	665.00	700.00	715.00	765.00	196.50
	23	731.00	790.00	\$45.00	900.00	935.00	960.00	990.00	259.00
	30	442.50	475.00	500.00	505.00	520.00	550.00	550.00	107.50
	43	614.00	640.00	670.00	725.00	750.00	790.00	830.00	216.00
Averag	e .	571.05	611.50	639.50	685.50	713.00	738.00	768.50	197.45

Lot No.	Animal No.	11-13-14 1947	12-11-47	1-8-48	2-5-48	3-4-48	4-8-48	5-5-6 1948	Total gain
II	5	614.00	665.00	690.00	725.00	745.00	785.00	815.00	201.00
	6	557.50	610.00	645.00	695.00	730.00	755.00	787.50	230.00
	18	677.00	710.00	740.00	815.00	\$40.00	845.00	900.00	223.00
	21	660.00	700.00	750.00	805.00	\$50.00	\$90.00	915.00	255.00
	24	553.00	565.00	610.00	640.00	660.00	680.00	717.50	164.50
	25	690.00	730.00	775.00	\$00.00	\$15.00	\$45.00	885.00	195.00
	33	581.50 -	565.00	610.00	650.00	680.00	695.00	737.50	156.00
	42	495.00	565.00	580.00	625.00	640.00	680.00	697.50	202.50
	39	648.00	695.00	725.00	775.00	800.00	\$30.00	\$55.00	207.00
	40	418.50	465.00	490.00	530.00	550.00	565.00	592.50	174.00
Average		589.45	627.00	661.50	706.00	731.00	757.00	790.25	200.80

D

Lot No.	Animal No.	11-13-14 1947	12-11-47	1-8-48	2-5-48	3-4-48	4-8-48	5-5-6 1948	Total gain
III	1	699.00	730.00	770.00	\$10.00	845.00	875.00	905.00	206.00
	11	556.00	595.00	635.00	690.00	710.00	725.00	777.50	221.50
	12	537.00	595.00	620.00	670.00	700.00	720.00	757.50	220.50
	13	727.50	760.00	775.00	\$30.00	840.00	865.00	900.00	172.50
	15	506.00	560.00	585.00	625.00	640.00	660.00	692.50	186.50
	27	540.00	570.00	615.00	650.00	680.00	710.00	757.50	217.50
	29	597.00	655.00	695.00	745.00	770.00	800.00	\$50.00	253.00
	31	582.50	630.00	660.00	700.00	730.00	745.00	787.50	205.00
	36	640.00	670.00	700.00	745.00	755.00	770.00	\$15.00	175.00
	37	570.00	580.00	610.00	665.00	680.00	705.00	712.50	142.50
Averag	e	595.50	634.50	666.50	713.00	735.00	757.50	795.50	200.00

SU

Lot No.	Animal No.	11-13-14 1947	12-11-47	1-8-48	2-5-48	3-4-48	4-8-48	5-5-6 1948	Total gain
IV	4	590.00	570.00	605.00	665.00	660.00	640.00	690.00	100.00
	8	565.00	570.00	605.00	665.00	660.00	660.00	695.00	130.00
	16	747.50	775.00	805.00	\$75.00	\$65.00	865.00	907.50	160.00
	19	560.00	580.00	620.00	650.00	655.00	660.00	695.00	135.00
	22	635.00	635.00	660.00	700.00	680.00	715.00	742.50	107.50
	26	536.00	550.00	580.00	630.00	640.00	630.00	672.50	136.50
	28	414.00	440.00	470.00	490.00	480.00	500.00	535.00	121.00
	32	637.00	675.00	705.00	720.00	725.00	730.00	742.50	105.50
	34	606.50	620.00	655.00	695.00	690.00	705.00	727.50	121.00
Averag	e	587.88	601.67	633.89	676.67	672.78	678.33	711.94	124.06

Lot No.	Animal No.	5-5-6 1948	6-5-48	7-8-48	8-5-48	9-2-48	9-30-48	11-4-5 1948	Total gain
I	2	970.00	970.00	1000.00	1020.00	1060.00	1095.00	1092.50	122.50
	3	\$65.00	815.00	\$70.00	\$70.00	915.00	960.00	935.00	70.00
	7	730.00	720.00	750.00	800.00	\$40.00	860.00	\$62.50	132.50
	9	780.00	750.00	\$10.00	\$40.00	880.00	925.00	935.00	155.00
	10	627.50	605.00	670.00	700.00	735.00	\$05.00	\$22.50	195.00
	14	575.00	555.00	610.00	640.00	660.00	700.00	717.50	142.50
	17	765.00	770.00	\$00.00	\$30.00	\$50.00	\$90.00	900.00	135.00
	23	990.00	950.00	980.00	1030.00	1010.00	1080.00	1025.00	35.00
	30	550.00	490.00	550.00	570.00	600.00	625.00	602.50	52.50
	43	830.00	780.00	840.00	850.00	895.00	920.00	932.50	102.50
Averag	ge	768.50	740.50	788.00	\$15.00	865.25	886.00	882.50	114.00

Lot No.	Animal No.	5-5-6 1948	6-5-48	7-8-48	8-5-48	9-2-48	9-30-48	11-4-5 1948	Total gain
II	5	815.00	\$20.00	\$50.00	880.00	890.00	945.00	922.50	107.50
	6	787.50	760.00	800.00	\$30.00	\$70.00	960.00	922.50	135.00
	18	900.00	885.00	910.00	920.00	980.00	980.00	955.00	55.00
	21	915.00	850.00	920.00	950.00	1000.00	1075.00	1000.00	85.00
	24	717.50	730.00	750.00	780.00	785.00	\$50.00	875.00	157.50
	25	\$85.00	875.00	\$70.00	910.00	950.00	990.00	1007.50	122.50
	33	737.50	755.00	800.00	\$40.00	850.00	900.00	\$70.00	132.50
	42	697.50	705.00	735.00	770.00	770.00	845.00	\$15.00	117.50
	39	855.00	\$25.00	850.00	\$\$0.00	920.00	965.00	977.50	122.50
	40	592,50	595.00	660.00	700.00	700.00	780.00	832.50	140.00
Averag	e	790.25	.780.00	814.50	846.00	871.50	929.00	907.75	117.50

Lot No.	Animal No.	5-5-6 1948	6-5-48	7-8-48	8-5-48	9-2-48	9-30-48	11-4-5 1948	Total gain
III	1	905.00	885.00	890.00	950.00	930.00	1050.00	1017.50	112.50
	11	777.50	785.00	790.00	800.00	\$55.00	885.00	895.00	117.50
	12	757.50	765.00	800.00	\$20.00	\$90.00	905.00	907.50	150.00
	13	900.00	880.00	900.00	920.00	950.00	980.00	950.00	50.00
	15	692.50	650.00	710.00	720.00	730.00	765.00	\$20.00	127.50
	27	757.50	740.00	770.00	790.00	830.00	865.00	885.00	127.50
	29	\$50.00	865.00	860.00	880.00	920.00	930.00	910.00	60.00
	31	787.50	770.00	\$30.00	\$50.00	\$35.00	895.00	887.50	100.00
	36	815.00	815.00	\$60.00	\$40.00	890.00	930.00	887.50	72.50
	37	712.50	705.00	760.00	790.00	\$10.00	\$35.00	832.50	120.00
Averag	;e	795.50	786.00	817.00	836.00	864.00	904.00	899.25	103.75

Lot No.	Animal No.	5-5-6 1948	6-5-48	7-8-48	8-5-48	9-2-48	9-30-48	11-4-5 1948	Total gain
IV	4	690.00	740.00	750.00	780.00	785.00	\$60.00	832.50	142.50
	8	695.00	715.00	760.00	800.00	\$25.00	865.00	857.50	162.50
	16	907.50	995.00	1020.00	1060.00	1080.00	1135.00	1127.50	220.00
	19	695.00	775.00	780.00	\$30.00	\$65.00	895.00	900.00	205.00
	22	742.50	805.00	850.00	910.00	915.00	1005.00	990.00	247.50
	26	672.50	725.00	730.00	750.00	775.00	830.00	807.50	135.00
	28	535.00	595.00	610.00	670.00	720.00	745.00	720.00	185.00
	32	742.50	\$40.00	\$50.00	\$70.00	885.00	865.00	917.50	175.00
	34	727.50	780.00	800.00	\$30.00	\$50.00	905.00	882.50	155.00
Averag	e	711.94	774.44	794.44	833.33	855.55	900.50	892.77	180.83

N

	Date	December	12, 1946	May 15, 1947	Nov. 14, 1947	May 6	, 1948
10.	No.	rype	Condición	туре	rybe	туре	Condicion
I	2 3 7 9 10 14 17 23 30 43	Good - Medium Medium / Medium / Good Good Good Good - Good -	Good - Medium - Medium - Medium - Medium - Medium Good - Good - Medium ≠	Good Good Medium Medium Good Medium Good Medium Good	Choice - Choice - Good Medium Medium Medium Medium Choice - Medium - Good	Medium Good / Medium Good - Medium - Medium / Good Common / Good -	Medium / Good / Medium Good - Medium - Medium / Good Medium - Good
I	56 18 21 25 33 29 40	Good - Good - Medium - Good / Good - Good - Medium Medium Good - Good -	Medium - Medium - Good - Medium / Medium Medium Medium - Good Medium	Medium Good Good Medium Good Medium Medium Good Medium	Good - Good - Medium / Good / Medium / Good Medium - Good Medium -	Medium / Medium / Good / Medium / Good - Medium Common / Medium Common	Medium / Medium / Good - Choice - Medium Medium / Medium / Medium / Medium / Common /

Lot No.	Date Animal No.	December Type	12, 1946 Condition	May 15, 1947 Type	Nov. 14, 1947 Type	May 6 Type	, 1948 Condition
III	1 11 12 13 15 27 29 31 36 37	Good / Good - Medium Good - Good - Medium / Medium / Good Good -	Medium / Medium / Medium - Medium Medium Medium / Good - Good - Good -	Good Good Medium Good Medium Good Medium Good Good	Choice - Good - Medium Choice Good - Good Good - Good Medium <del>/</del> Good	Good Good Medium Good / Good - Good - Medium - Good - Good -	Good / Good / Medium Choice - Good Good Medium - Good Good -
IV	4 8 16 19 20 22 26 28 32 34	Good - Medium Good ≠ Medium Good - Good - Medium Good Good - Medium ≠	Medium Medium Good - Medium Medium - Medium - Good - Medium / Good -	Good Medium Good Medium Medium Good Medium Good Medium	Medium Good - Good / Good - Died November Good - Good Medium Good Medium	Medium / Good - Common / 11, 1947. Medium Good Common / Medium / Good -	Medium / Good - Medium - Medium - Medium / Medium / Good -

Table 20FEED COS	STS 1946-4	8		
	Lot I	Lot II	Lot III	Lot IV
Winter 1946-47 Chopped cane Soybean oil meal Alfalfa hay Fish oil Mineral mixture Salt Pasture Total	\$101.34 65.33  1.85 0.80 \$169.32	\$101.79 65.33 0.94 1.85 1.00 	\$ 82.38 43.50 38.44  2.27 1.30 \$167.89	\$ 3.00 64.90  0.21 0.20 55.00 \$123.31
Av/head	16.93	17.09	16.79	12.33
Summer 1947 Pasture Av/head	\$ 75.00 \$ 7.50	\$ 75.00 \$ 7.50	\$ 75.00 \$ 7.50	\$ 75.00* \$ 7.50
Winter 1947-48 Chopped cane Soybean oil meal Alfalfa hay Fish oil Mineral mixture Salt Pasture	\$279.80 91.35  1.40 1.20	\$281.21 91.35 1.17 1.60 1.50	\$261.49 56.31 32.88  1.40 1.60	<pre>\$ 10.80 81.74 0.60 0.70 67.50</pre>
Total	\$373.75	\$376.83	\$353.68	\$161.34
Av/head	37.38	37,68	35.37	17.93
Summer 1948 Pasture Av/head	\$120.00 \$ 12.00	\$120.00 \$ 12.00	\$120.00 \$ 12.00	\$108.00
Grand Total Av/head	\$738.07 73.81	\$742.74 74.27	\$716.57 71.66	\$467.65
* Heifer number	20 died N	ovember 1	1, 1947.	
Feed costs used: Chopped cane Alfalfa hay Soybean oil meal Fish liver oil Mineral mixture Salt Pasture - Head/mo	Ton Ton Ton Lb. Cwt. Cwt. onth	Winter S 1946-7 12.50 25.00 115.00 1.62 41.20 20.00 1.00	ummer       Wintender         1947       1947         1947       1947         25       25         105       105         105       1         40       20         11.25       1	Summer -8 1948 .00 .00 .00 .00 .00 .00 .25 \$2.00



## BIBLIOGRAPHY

- Boyer, P. D., Phillips, P. H., Lundquist, N. S., Jensen, C. W., and Rupel, I. W. Vitamin A and carotene requirements for the maintenance of adequate blood plasma vitamin A in the dairy calf. Journal of Dairy Science, 25:433-440, May 1942.
- Boyer, P. D., Phillips, P. H., and Smith, J. K. The separation of carotene from vitamin A for the determination of vitamin A in blood plasma. Journal of Biological Chemistry, 152:445-452, February 1944.
- 3. Braun, Werner. Studies on the carotenoid and vitamin A levels in cattle. I. Seasonal changes of the carotenoid and vitamin A levels and the normal carotenoid-vitamin A ratio of the blood. Journal of Nutrition, 29:61-71, January 1945.
- 4. Braun, Werner. Studies on the carotenoid and vitamin A levels in cattle. II. Carotenoids and vitamin A in the liver, their ratio and their relationship to blood levels. Journal of Nutrition, 29:73-79, January 1945.
- Briggs, A. P. A modification of the Bell-Doisy phosphate method. Journal of Biological Chemistry, 53:13-16, May 1922.
- 6. Brody, S. Bioenergetics and growth. New York, Reinhold Publishing Corporation, 1945. 1023 p.
- 7. Clark, E. P., and Collip, J. B. A study of the Tisdall method for the determination of blood serum calcium with a suggested modification. Journal of Biological Chemistry, 63:461-464, March 1925.
- Converse, H. T., and Meigs, Edward B. Some disasters in reproduction and growth caused by low quality hay. American Society of Animal Production. Proceedings, 24:141-144, 1932.

 Converse, H. T., and Meigs, Edward B. The carotene requirements for normal reproduction. Journal of Dairy Science, 21:114-115, May 1938.

- Converse, H. T., and Meigs, Edward B. The vitamin A requirement for normal growth in young dairy cattle. American Society of Animal Production. Proceedings, 32:67-72, January 1939.
- 11. Crowther, Charles. Nutrition and reporduction. East African Agricultural. J. Kenya, Tanganyika, Uganda, Zanzibar, 4:327-333, 1939. Abstracted in Chemical Abstracts, 35:3296, May 20, 1941.
- Dann, W. J., and Evelyn, K. A. Determination of vitamin A with the photoelectric colorimeter. Biochemical Journal, 32:1008-1017, June 1938.
- 13. Davis, Russell E., and Madsen, Louis L. Carotene and vitamin A in cattle blood plasma with observations on reproductive performance at restricted levels of carotene intake. Journal of Nutrition, 21:135-146, February 1941.
- Dickson, R. E., and Jones, J. H. Vitamin A in the cattle wintering ration in Texas. Texas Agricultural Experiment Station, Annual Report 48: 200-201, 1935.
- 15. Fernandes, C. J. Famine and vitamin A deficiency in cattle. Indian Farming, 1(12):591-592. Abstracted in Biological Abstracts, 19:6642, April 1945.
- 16. Flora, C. C., Ward, R. E., Bechdel, S. I., Guerrant, N. B., and Dutcher, R. A. Carotene in the nutrition of dairy calves. Journal of Dairy Science, 22:321-328, May 1939.
- 17. Gallup, Willis D., and Kuhlman, A. H. Carotene content of the blood plasma of dairy cattle in relation to vitamin A deficiency. Oklahoma Academy of Science. Proceedings, 21:89-92, 1941. Abstracted in Biological Abstracts, 16:543, January 1942.
- 18. Garrigus, W. P., and Rusk, H. P. Some effects of the species and stage of maturity of plants on the forage consumption of grazing steers of various weights. Illinois Agricultural Experiment Station. Bulletin, 454:441-508, 1939.

19. Guilbert, H. R. and Hart, G. H. Minimum vitamin A requirement with particular reference to cattle. Journal of Nutrition, 10:409-427, October 1935. 20. Guilbert, H. R. and Hart, G. H. Storage of vitamin A in cattle. Journal of Nutrition, 8:25-44, July 1934. 21. Guilbert, H. R., Howell, C. E., and Hart, G. H. Minimum vitamin A and carotene requirements of mammalian species. Journal of Nutrition, 19:91-103, January 1940. 22. Guilbert, H. R., Miller, R. F., and Hughes, E. H. The minimum vitamin A and carotene requirements of cattle, sheep and swine. Journal of Nutrition, 13:543-564, May 1937. 23. Hansen, R. Gourth, Phillips, Paul H., and Rupel, I.W. The effect of vitamin supplements on survival of newborn calves. Journal of Dairy Science, 29:761-766, November 1941. 24. Hart, G. H. and Guilbert, H. R. Vitamin A deficiency as related to reproduction in range cattle. California Agricultural Experiment Station. Bulletin, 560:1-30, 1933. Hart, G. H., Guilbert, H. R., and Goss, H. Seasonal 25. changes in the chemical composition of range forage and their relation to nutrition of animals. California Agricultural Experiment Station. Bulletin, 543:3-62, 1932. 26. Hart, G. H., Mead, S. W., and Guilbert, H. R. Vitamin A deficiency in cattle under natural conditions. Society for Experimental Biology

- conditions. Society for Experimental Biology and Medicine. Proceedings, 30:1230-1233, 1933. Abstracted in Nutrition Abstracts and Reviews 3:1542, October 1933.
- 27. Jones, J. H., Riggs, J. K., Fraps, J. S., Jones, J. M., Schmidt, H., Dickson, R. E., Howe, Paul E., and Black, W. H. Carotene requirements for fattening beef cattle. American Society of Animal Production. Proceedings, 31:94-102, 1938.

	128
28.	Jones, J. H., Schmidt, H., Dickson, R. E., Fraps, G. S., Jones, J. M., Riggs, J. K., Kemmerer, A. R., Howe, P. E., Black, W. H., Ellis, N. R., and Marion, Paul T. Vitamin A studies in fattening feeder calves and yearlings. Texas Agricultural Experiment Station. Bulletin, 630:1-52, 1943.
29.	Keener, H. A., Bechdel, S. I., Guerrant, N. B., and Thorp, W. T. S. Carotene in calf nutrition. Journal of Dairy Science, 25:571-583, July 1942.
30.	Kempt, W. B. A summary of wartime research. Mary- land Agricultural Experiment Station. Annual Report, 57:1-168, 1944.
31.	Kimble, M. S. The photocolorimetric determination of vitamin A and carotene in human plasma. Journal of Laboratory and Clinical Medicine, 24:1055, 1939.
32.	Kniese, Marvin E. Alfalfa hay supplement as com- pared to fish liver oil in the vitamin A nutrition of eastern Colorado beef cattle. Master's thesis, 1947. Colorado A & M College. 68 p. ms.
33.	Kuhlman, A. H. and Gallup, W. D. Carotene (pro- vitamin A) requirements of dairy cattle for conception. Journal of Dairy Science, 25: 688-689, August 1942.
34.	Kuhlman, A. H. and Gallup, W. D. Carotene (pro- vitamin A) requirement of dairy cattle for reproduction. American Society of Animal Production. Proceedings, 33:67-73, December 1940.
35.	Kuhlman, A. H. <u>and</u> Gallup, W. D. The relation of carotene intake to blood carotene values of dairy cattle. Journal of Animal Science, 1:68, February 1942.
36.	Lewis, J. M. <u>and</u> Wilson, Logan T. Vitamin A requirements in calves. Journal of Nutrition, 30:467-475, December 1945.
37.	Madsen, Louis L., Earle, I. P., Davis, Russell E., Bastron, Harry, Thies, C. W., Cabell, Charles A., and Ray, H. Dean. Some observations on beef cattle affected with generalized edema or anasarca due to vitamin A deficiency. Journal of Nutrition, 34:603-619, December 1947.

	129
38.	Madsen, Louis L., Eaton, C. N., Heemstra, Louis, Davis, Russell E., Cabell, Charles A., <u>and</u> Knapp, Bradford Jr. Effectiveness of caro- tene and failure of ascorbic acid to increase sexual activity and semen quality of vitamin A deficient beef bulls. Journal of Animal Science, 7:60-69, February 1948.
39.	Maynard L. A. Animal nutrition; 2d. ed. New York, McGraw-Hill Book Company, Inc., 1947. 494 p.
40.	Moore, L. A. Carotene intake, level of blood plasma carotene, and the development of papil- lary edema and nyctalopia in calves. Journal of Dairy Science, 22:803-812, October 1939.
41.	Moore, L. A. Relationship between carotene, blind- ness due to constriction of the optic nerve, papillary edema and nyctalopia. Journal of Nutrition, 17:443-459, May 1939.
42.	Moore, L. A. The effect of pasture on the carotene content of blood plasma of the bovine. Jour- nal of Dairy Science, 22:513-520, July 1939.
43.	Moore, L. A. Vitamin A in reproduction and milk production. American Miller and Processor, 72(12):60-62, 1944. Abstracted in Biological Abstracts, 19:8720, May 1945.
44.	Moore, L. A., <u>and</u> Berry, M. H. Vitamin A and carotene content of the blood plasma of dairy calves from birth up to four months of age. Journal of Dairy Science, 28:821-826, November 1945.
45.	Moore, L. A., Berry, M. H., <u>and</u> Sykes, J. F. Caro- tene requirements for the maintenance of a normal spinal fluid pressure in dairy calves. Journal of Nutrition, 26:649-658, December 1943.
46.	Morrison, F. B. Feeds and feeding; 20th edition. Ithaca, New York, The Morrison Publishing Company, 1941. 1050 p.
47.	Nelson, H. F., Moore, L. A., Horwood, R. E., and Branaman, G. A. Vitamin A and carotene con- tent of the blood plasma of beef and dairy calves from birth to four months of age. Michigan Agricultural Experiment Station. Quarterly Bulletin, 27(1):50-53, August 1944.

- 48. Payne, Merle G., <u>and</u> Kingman, H. E. Carotene blood levels and reproductive performance in range Hereford cattle. Journal of Animal Science, 6:50-55, February 1947.
- 49. Peterson, W. J. Recent developments in methods for determining carotene. Industrial and Engineering Chemistry, Analytical Edition, 13:212-216, April 15, 1941.
- 50. Riggs, J. K. Length of time required for depletion of vitamin A reserves in range cattle. Journal of Nutrition, 20:491-500, November 1940.
- 51. Sampson, J., Boley, L. E., <u>and</u> Graham, Robert. Losses of newborn calves presumably the result of feeding dams rations inadequate for reproduction. Cornell Veterinarian, 28:53-57, January 1938.
- 52. Schmidt, H. Mineral requirements of cattle in the Red Beds region of Texas. Texas Agricultural Experiment Station. Annual Report, 48:19-20, 1935.
- 53. Sutton, T. S., and Soldner, P. A. Seasonal variation in the blood plasma carotene and vitamin A of adult dairy cattle. Journal of Dairy Science, 28:859-867, November 1945.
- 54. Ward, R. E., Bechdel, S. I., <u>and</u> Guerrant, N. B. Carotene and vitamin A in the nutrition of growing dairy cattle. Journal of Dairy Science, 23:115-124, February 1940.
- 55. Washburn, L. E., Connell, W. E., and Wheeler, S. S. Do vitamin A supplements need to be added to beef cattle feed in Colorado? Colorado A & M News, 1:6, December 1946.

LIBRARY GOLORADO A. & M. COLLEGE FORT COLLINS COLORADO