

THESIS

CLIMATIC AND HYDROLOGIC PROCESSES LEADING TO RECENT WETLAND LOSSES IN
YELLOWSTONE NATIONAL PARK, USA

Submitted by

Derek M. Schook

Graduate Degree Program in Ecology

In partial fulfillment of the requirements

For the Degree of Master of Science

Colorado State University

Fort Collins, Colorado

Spring 2012

Master's Committee:

Advisor: David J. Cooper

Michael J. Ronayne
Stephanie K. Kampf

ABSTRACT

CLIMATIC AND HYDROLOGIC PROCESSES LEADING TO RECENT WETLAND LOSSES IN YELLOWSTONE NATIONAL PARK, USA

Wetlands provide vital habitat within functioning environments and act as landscape indicators by integrating catchment-scale hydrologic processes. Wetland drying during the past few decades in Yellowstone National Park's Northern Range has caused concern among National Park managers and the public at large. My research was initiated to develop an understanding of the processes controlling wetland water levels and contributing to wetland decline in the Northern Range. To do this I integrated analyses of hydrology, climate, soils, and vegetation. In 2009 I selected 24 study wetlands and instrumented each with an average of five shallow groundwater monitoring well-and-piezometer nests. To quantify historic wetland area I mapped hydric soils, analyzed aerial photographs, and identified geomorphic indicators of higher water. Vegetation was sampled to characterize wetlands and plant-water relationships, and I also conducted a soil seed bank study. The Trumpeter Lake focal site revealed groundwater changes through time and was used to identify the timescale on which an important wetland varies. Climate data indicated that warming and drying occurred during the 20th century, but that this pattern was within the natural range of variation for the study region during the past 800 years. Hydrologic data revealed that study sites included locations of groundwater discharge, recharge, and flow-through as well as water perched above the regional water table. Hydrologic regimes were classified using a shape-magnitude framework and seven wetland classes were characterized. Wetland classes exhibited variable hydrologic permanence within and between the two study summers. Aerial photographs and hydric soil delineation both confirmed formerly greater wetland abundance. These changes were linked to the wetland classes and the presence or absence of surface water outlets. Wetland plant species inhabited areas of distinct water

table depth and variation, and can be used to infer subsurface hydrologic regime in the absence of extensive monitoring well networks. Continued monitoring of these wetland basins and their watersheds is critical to expanding our understanding of the processes supporting Northern Range wetlands and allowing us to better manage these valuable habitats.

ACKNOWLEDGEMENTS

This work to which I sign my name as sole author is the culmination of efforts by many people. My advisor, Dr. David Cooper, has been instrumental to the project's success from day one, guiding research objectives and my personal and professional development throughout. The project was funded through grants from the Canon Foundation, Yellowstone Foundation, National Park Service, and Warner College of Natural Resources. Roy Renkin's contributions aided field work and facilitated the CSU-NPS collaboration. Mary Hektner's role as park liaison was valuable in developing research objectives and understanding NPS goals. Thanks to collaboration with the Yellowstone Center for Resources, including Christie Hendrix, Stacey Gunther, Hank Heasler, Cheryl Jaworowski, David Susong, and Tom Olliff. I thank Erin Ouzts for a fun summer of dedicated field work. Jennifer Whipple identified dozens of wetland plant species when my efforts came up short. Special thanks to my colleagues in the Cooper lab for mulling over the project's details, reviewing drafts, and engendering an enjoyable work environment. Stephen Gray shared climate reconstruction data, and Ken Pierce offered seasoned opinions of Northern Range geology and hydrology. Committee members Mike Ronayne and Stephanie Kampf supplied insight regarding the scope and goals of the project, as well as helpful revisions. This project and my graduate degree have been facilitated by the Graduate Degree Program in Ecology and the department of Forest and Rangeland Stewardship at Colorado State University.

TABLE OF CONTENTS

1. INTRODUCTION	1
2. STUDY AREA	3
3. METHODS	5
3.1 WETLAND CHARACTERISTICS AND CLASSIFICATION	5
<i>3.1.1 Hydrologic Data collection</i>	<i>5</i>
<i>3.1.2 Wetland Water Table Classification</i>	<i>6</i>
<i>3.1.3 Wetland area analysis</i>	<i>8</i>
3.2 CLIMATE	9
<i>3.2.1 2009-2010 weather</i>	<i>9</i>
<i>3.2.2 Historic datasets</i>	<i>10</i>
3.3 VEGETATION	10
<i>3.3.1 Plant-hydrology relationship</i>	<i>10</i>
<i>3.3.2 Soil seed bank study</i>	<i>11</i>
3.4 TRUMPETER LAKE FOCAL SITE	12
<i>3.4.1 Surveying and aerial photograph analysis</i>	<i>12</i>
<i>3.4.2 Aquifer flow paths</i>	<i>13</i>
4. RESULTS	13
4.1 WETLAND HYDROLOGIC REGIMES	13
<i>4.1.2 Wetland Classification</i>	<i>16</i>
<i>4.1.3 Wetland size through time</i>	<i>18</i>
4.2 CLIMATE	19
<i>4.2.1 Trends in Yellowstone River discharge</i>	<i>19</i>
<i>4.2.2 Long-term climate trends</i>	<i>20</i>
4.3 VEGETATION	20
<i>4.3.1 Vegetation-hydrology relationship</i>	<i>20</i>
<i>4.3.2 Soil seed bank study</i>	<i>21</i>
5. DISCUSSION	22
5.1 HYDROLOGICALLY-BASED WETLAND CLASSIFICATION	23
5.2 HYDROLOGIC PROCESSES CONTROLLING YELLOWSTONE’S WETLANDS	24
5.3 HYDROLOGIC CONTROLS OF WETLAND VEGETATION	24
5.4 TRUMPETER LAKE WATERSHED DYNAMICS	25
5.5 WEATHER, CLIMATE, AND WETLAND AREA CHANGES THROUGH TIME	27
6. CONCLUSIONS	29
7. FIGURES AND TABLES	31
8. REFERENCES	47
9. APPENDICES	54
APPENDIX A. ADDENDUM TO 2009-2010 WEATHER REPORT	54
APPENDIX B. LOCATION OF ALL WELLS	56
APPENDIX C. AERIAL PHOTOGRAPHY DATES	60
APPENDIX D. HYDRAULIC CONDUCTIVITY MEASUREMENTS IN THE TRUMPETER LAKE WATERSHED	61
APPENDIX E. TRUMPETER LAKE FOCAL SITE ANALYSIS	63

APPENDIX F. SOIL SEED BANK STUDY SPECIES LIST	65
APPENDIX G. SOIL SEED BANK STUDY PLOT STRATIFICATION PROCEDURE	67
APPENDIX H. EXAMPLE WETLAND HYDROGRAPHS FROM THE SEVEN CLASSES	70
APPENDIX I. YELLOWSTONE RIVER ANNUAL DISCHARGE.....	72
APPENDIX G. PHOTOGRAPHS OF ONE WETLAND FROM EACH CLASS	73

LIST OF FIGURES

FIGURE 1. MAP OF THE YELLOWSTONE NATIONAL PARK AND ITS NORTHERN RANGE, USA	31
FIGURE 2. CLUSTER ANALYSIS DENDROGRAMS OF WATER TABLE VARIATIONS IN WELLS	32
FIGURE 3. HYDROGRAPHS OF THE SEVEN WELL CLASSES.	33
FIGURE 4. CROSS SECTIONS AND HYDROGRAPHS FOR TWO EXAMPLE WETLANDS	35
FIGURE 5. CROSS SECTION OF TRUMPETER LAKE WITH WATER TABLES FROM FOUR DATES.....	36
FIGURE 6. TRUMPETER LAKE WATER TABLE CONTOURS AND LAKE STAGES, 2010.....	37
FIGURE 7. SURFACE AREA-VOLUME RELATIONSHIP AT TRUMPETER LAKE.	38
FIGURE 8. STUDY AREA CLIMATE RECONSTRUCTION	39
FIGURE 9. YELLOWSTONE RIVER DISCHARGE SHAPE-MAGNITUDE ANALYSIS CLASSES	40
FIGURE 10. COMPARISON OF EARLY AND LATE YELLOWSTONE RIVER RUNOFF YEARS	41
FIGURE 11. PLANT SPECIES WATER TABLE HYDROGRAPHS	42
FIGURE 12. WETLAND PLANTS GERMINATING IN THE SOIL SEED BANK STUDY.....	43

LIST OF TABLES

TABLE 1. DESCRIPTION OF THE 24 WETLAND STUDY SITES 44

TABLE 2. ENVIRONMENTAL INFLUENCES ON WETLAND CLASS ASSIGNMENT 45

TABLE 3. TEMPORAL RELATIONSHIP BETWEEN PRECIPITATION ON AND WETLAND SIZE..... 46

TABLE 4. TRUMPETER LAKE SIZE RELATED TO PRECIPITATION FROM THE LAST DECADE..... 46

1. INTRODUCTION

Wetlands are among the most valuable yet vulnerable habitats on Earth (Poff et al., 2002; Bates et al., 2008; Winter, 2000). Hydrologic processes, which are influenced by climate, geology, and landscape setting, are the dominant mechanisms creating and sustaining wetlands (Mitsch and Gosselink, 2000; Hunt et al., 1996). Located at low points in their watersheds, wetlands integrate catchment-scale processes and reflect environmental conditions (Williamson et al., 2008; Bates et al., 2008; Long and Nestler, 1996). Additionally, the close proximity of the water table and land surface in wetlands leaves these habitats susceptible to changing hydrologic, landscape, and climatic conditions (Brooks, 2009; Bates et al., 2008).

Interactions among precipitation, evapotranspiration, groundwater, and surface water create a wetland's hydrologic regime. Distinguishing surface- from groundwater processes is often complex because they interact at multiple spatial and temporal scales (Winter, 1999; Schot and Winter, 2006; Devito et al., 2005). Subsurface stratigraphy and its hydraulic properties play important roles in influencing wetland hydrologic processes (van der Kamp and Hayashi, 2009; Winter, 1999; Todd et al., 2006). Adjacent wetlands that appear similar can vary in permanence (Brooks, 2000), and recent modeling in the prairie pothole region of the northern American Great Plains has shown more permanent wetlands to be more susceptible to climate change than temporary wetlands (Johnson et al., 2010). Further complicating hydrologic assessments, groundwater sources sustaining a wetland may originate hundreds of kilometers away, such as those that support Argentine desert oases (Jobbágy et al., 2011) and Death Valley, California springs (Belcher et al., 2009). Groundwater flow paths also affect surface water chemistry (van der Kamp and Hayashi, 2009; LaBaugh, 1987) and salinity (Jolly et al., 2008; LaBaugh et al., 1998), factors that influence wetland species composition.

Basin wetlands found throughout the world exhibit a range of hydrologic regimes (Winter, 1999). For example, playa wetlands of the southern Great Plains (Tiner, 2003; Tsai et al., 2007) and vernal pools across the USA (Brooks, 2004; Pyke, 2004; Zedler, 2003) are dependent on precipitation and evapotranspiration (ET) processes. These wetlands are considered hydrologically isolated from ground- and surface waters. In contrast, Nebraska's Sandhills and Colorado's Great Sand Dunes support wetland complexes that are connected via groundwater flows in highly conductive soil (Winter, 1986; Wurster et al., 2003). Cook and Hauer (2007) described wetlands in the Northern Rocky Mountains that formed in a dead-ice glacial moraine, with some wetlands connected by near-surface flow and others being hydrologically isolated. Fluxes between ground- and surface water in prairie pothole region wetlands are highly variable temporally and spatially, and the direction of groundwater flow may change seasonally (Woo and Rowsell, 1993; Rosenberry and Winter, 1997; Winter and Rosenberry, 1995). The high degree of spatial and temporal variability in wetlands illustrates that generalizing wetland function in an unstudied region can yield inaccurate assumptions.

Temperature and precipitation strongly influence wetland habitats and are forecasted to change in the coming decades (Carpenter et al., 1992; Brooks, 2009). Past changes in climate have been correlated with wetland disappearance in Alaska and Siberia (Klein et al., 2005; Smith et al., 2005), and the trend is predicted to continue (Sorenson et al., 1998; Bates et al., 2008). Understanding the effects of climate change on wetland biotic and hydrologic processes is challenging due to the inherent spatial and temporal complexity in these habitats (Pilon and Yue, 2002; Bates et al., 2008; Brooks, 2009).

Regional climate models for Yellowstone National Park (YNP) forecast an ecological shift unprecedented in the Quaternary, yielding an uncertain future for the park's water-based ecosystems (Westerling et al., 2011; Bartlein et al., 1997). Many wetlands in YNP's glacially-

influenced Northern Range have exhibited pronounced surface water declines during the past four decades. However, these changes are poorly quantified (e.g., McMenamin et al., 2008), and no previous research has investigated the processes causing these changes. Recent wetland loss has already affected YNP's native wetland species, including causing the loss of trumpeter swan nesting habitat (Proffitt et al., 2010). Although some effects of wetland decline on key species have been identified, the underlying processes altering the wetlands themselves remain unknown and are important to guiding future conservation efforts.

The geologic diversity of YNP has created high environmental heterogeneity, precluding a broad characterization of wetland processes. To address this variation, I used concepts from other classifications (e.g., Acreman et al., 2009; Dahl, 2011; Rains, 2011; Junk et al., 2011) to create a framework for grouping wetlands according to their hydrologic processes. This research addressing wetland change was framed within the following objectives: (1) classify wetlands according to their hydrologic regime, including climatic and geomorphic processes, (2) determine the patterns and magnitude of water level decline that occurred during the late 20th and early 21st centuries and assess whether this change is within the natural range of variation, and (3) investigate a focal site that has experienced dramatic water level changes to gain a more complete understanding of wetland processes. The upper Yellowstone River watershed provides a unique opportunity to investigate hydrologic processes in a relatively pristine hydrologic setting. To address the study objectives I integrated wetland and watershed hydrology, climate, soils, and vegetation to create an integrated view of the processes supporting Northern Range wetlands.

2. STUDY AREA

2.1 Site Description

The 1,400 km² Northern Range is located in Wyoming and Montana and comprises much of YNP's northern quarter (Fig. 1). Douglas fir (*Pseudotsuga menziesii* Mirbel) and lodgepole pine (*Pinus contorta* Loudon) dominate the higher elevation forests, while lower elevations are dominated by sagebrush steppe (*Artemisia tridentata* Nutt; nomenclature follows USDA PLANTS (NRCS, 2011)). Lower elevations (1800-2000 m) receive 41 cm average annual precipitation, with over half falling as snow (NCDC, 2011). Most of the study area was covered by glaciers during the Pinedale Glaciation, which ended approximately 15 kya. Today's environment is heavily influenced by glacial scouring and till deposition that has created a heterogeneous landscape with abundant depressions that support today's wetlands. Clay rich mollisols and inceptisols surround the study sites (YCR, 2009). In recent years, park staff and visitors have observed a pronounced lowering of surface water levels in these wetlands, but quantitative and process-based information explaining the phenomenon is lacking.

In 2009, 24 non-riparian wetlands were selected as study sites to characterize Northern Range wetland types (Table 1). Study sites ranged from 1783 – 2284 m elevation. Sites included both mineral and organic soils, and common plant species included *Carex atherodes*, *C. utriculata*, *Juncus arcticus*, *Eleocharis palustris* and *Schoenoplectus acutus*. Study wetlands receive water from direct precipitation, groundwater, and overland flows. Through the 20th and early 21st centuries, some Northern Range wetland water levels were stable, while others varied greatly (Engstrom et al., 1991). Several wetlands exhibited distinct indicators of former high water levels, including relict vegetation, lichen trimlines, and eroded shorelines. Biotic factors including high large herbivore density can disturb vegetation and soils, especially in wetlands. Northern Range bison populations are the highest on record and elk populations have been nearly twice as high during the past three decades compared to the previous four (YNP, 1997; Wallen, 2010; Wyman, 2011).

The Trumpeter Lake focal site was selected for a more detailed analysis because it has purportedly undergone major water level declines and trumpeter swans formerly nested here. However, lower water levels have changed the lake's habitat structure, leaving it unsuitable for nesting today. The lake is located near the confluence of the Lamar and Yellowstone Rivers in dead-ice moraine terrain. Its watershed has hummocky topography comprised of low-permeability unconsolidated glacial till with a high density of granitic glacial erratics (Pierce, 1979). Upland soils were identified as loam and lake-bottom soil as clay-loam using the hydrometer method particle size analysis (Gee and Bauder, 1979), suggesting that porosities were approximately 46% (Rawls et al., 1982).

2.2 Study Period weather

Weather patterns during the study period influenced the measured wetland water levels. 2009 and 2010 annual temperatures were both within 0.3° C of the post-1931 average. Total precipitation in water year 2009 (1 Oct 2008 – 30 Sep 2009) was 97% and snowfall 120% of average, while 2010 total precipitation was 83% and snowfall 53% of average (NCDC, 2011). Early June through early July was the wettest period in 2009, while late May through mid-June was the wettest period in 2010 (Appendix A). Weather station comparison for 2010 data indicated that sites throughout the study area experienced similar precipitation conditions. Pearson's R correlation coefficients among the three rain gauges within the study area ranged from 0.84 to 0.97.

3. METHODS

3.1 WETLAND CHARACTERISTICS AND CLASSIFICATION

3.1.1 Hydrologic Data collection

103 shallow groundwater monitoring wells were installed at the 24 wetland study sites in 2009, and 18 wells were added in 2010 (Appendix B). Wells were distributed to measure water tables on all sides of each study wetland and to characterize the hydrologic niches of dominant plant species. Staff gauges were installed in all wetlands with ponded surface water to measure its depth. Monitoring wells were hand-augered using a 10 cm diameter bucket auger to a depth either below the anticipated water table low or as deep as possible in rocky soils. Wells were constructed from 4.2 cm I.D. schedule 40 PVC pipe that was continuously slotted throughout the anticipated zone of water table fluctuation. Augered holes were backfilled with native soil. To measure hydraulic head in various soil layers, an average of two nested 2.1 cm I.D. PVC piezometers were placed adjacent to each well within the top two meters of soil. Water depth was measured manually with an electric tape approximately biweekly in 2009 and weekly in 2010 (Shaffer et al., 2000), and readings were used to interpolate weekly values. A rotating laser level was used to determine the relative elevation of wells within a wetland.

3.1.2 Wetland Water Table Classification

Classifying monitoring wells based on measured water tables was the first step toward wetland classification. The majority of wells were installed in May 2009, and I used 3 June 2009 as the first date for analyzing well, staff gauge, and piezometer data. I excluded from analysis instruments installed in late 2009 or in 2010 along with those that dried too early to provide data throughout each summer. For classification, each well's initial water level was standardized to a common datum, and subsequent readings were relative to this point. Standardization permitted the ecologically significant analysis of surface and groundwater changes independent of their absolute ground-surface elevations (van der Kamp and Hayashi, 2009).

I grouped wells with similar water table variations using a hierarchical combination of multivariate statistics and well-nest informed assignment. Wells with two distinctive hydrologic regimes were first identified by analyzing hydrologic patterns. Wells in a stable group had less than 3 cm of water level variation during the two-year study period. Secondly, wells in a perched group had surface water that disappeared abruptly and nested piezometers that never contained water, revealing an unsaturated layer below surface water. These groups were deemed unique from all other wells, which contained transient water levels connected to groundwater systems.

For the remaining 83 wells, I conducted a two-step “shape-magnitude” cluster analysis to produce a composite classification by separately analyzing the timing (i.e. shape) and magnitude of water table variations. This approach has been used for analyzing weather patterns and stream discharge (Laize and Hannah, 2010; Bower et al., 2004; Hannah et al., 2000), and recently groundwater (Upton and Jackson, 2011). To classify shape, each well’s weekly water table data were standardized to a common degree of variation using a z-scores transformation (mean = 0, st. dev. = 1). This transformation isolated the seasonality and rate of water table change independent of its magnitude. Transformed data were grouped using hierarchical agglomerative cluster analysis using Euclidian distance and Ward’s group linkage method (Laize and Hannah, 2010; Bower et al., 2004) with the program PC-ORD (McCune and Mefford, 2006). The resultant dendrogram was pared at 40% information remaining, producing three groups of wells, each with distinct hydrograph characteristics.

For the magnitude analysis, I combined seven water table variables: minimums, means and standard deviations for each 2009 and 2010, and maximum elevation for 2010 (Bower et al. 2004, Harris et al. 2000). All magnitude indices were standardized by conversion to z-scores to eliminate uneven weighting of classes (Hannah et al., 2000; van Tongeren, 1987). A second

cluster analysis was performed on the magnitude data using the procedure described above but pruned at 0% information remaining, which produced two groups, one with larger and one with smaller water table changes (Fig. 2). The three shape and two magnitude classes were then crossed to yield six possible well hydrograph classes, five of which occurred in the study wetlands. The five shape-magnitude classes combined with the perched and stable well groups identified previously produced seven well classes (Fig. 3). Wells within most wetlands were in a single class, allowing wetlands to be grouped by these classes. A unique wetland class was created to accommodate the two sites that contained wells from three or four well classes.

To compare the wetland classification to local environments, I compared wetland classes to 14 environmental variables. Chi-squared analysis was conducted on the six binary categorical variables: surface water inflow, surface water outflow, peat presence, organic matter in basin, clay in basin, and bulrush (*Schoenoplectus acutus*) as the dominant wetland plant species. I define a wetland “basin” as a depression with surface water, not synonymous with a wetland’s watershed. One-way ANOVA, similar to chi-square analysis but used for quantitative variables, was conducted on eight variables: elevation, average annual precipitation, maximum observed surface water area, watershed size, duration to slowest piezometer’s equilibration, maximum piezometer positive head, electrical conductivity (EC) in the basin, and EC of groundwater inflow. To supplement these analyses, I created a classification tree via CART analysis (De’ath and Fabricius, 2000) in the rpart package (Therneau and Atkinson, 2009) using the statistical program R 2.10.1 (R Development Core Team, 2009).

3.1.3 Wetland area analysis

Hydric soils: At 16 of the 24 study sites the maximum elevation of wetland soils could be identified using the hydric soil indicators protocol in the Corps of Engineers’ Wetlands

Delineation Manual (USACE, 2010). The boundary between wetland and upland soil was determined using soil pits and morphological indicators including chroma less than two, oxidized root channels, and mottled matrices. The distance between modern water level and the hydric soil's upper boundary was analyzed among wetland classes using an ANOVA and between outlet vs. closed-basin wetlands using a t-test.

Aerial photograph analysis: Aerial photographs from 1954, 1969, 1991, 1994, 1998, 2001, 2006, and 2009 were used to quantify the ponded area through time (photo dates in Appendix C). Dense vegetation obscured the identification of surface water perimeters at ten of the study sites, so 14 wetlands were used in the analysis. Photos were georectified to 2009 NAIP imagery using 2nd and 3rd degree polynomials, and wetland surface area was delineated in ArcMap v. 10 (ESRI, 2010). To standardize wetland area (Niemuth et al., 2010), each wetland's maximum area was assigned 1, and other years represented a proportion of 1. Wetland areas were compared to annual precipitation totals (NCDC, 2011) at either the Tower or Mammoth weather station, depending on proximity. I used regression models between the two stations to estimate missing monthly precipitation values (Iglesias et al., 2006; Pegram and Pegram, 1993). SAS (SAS Institute, 2010) was used to conduct a multiple regression comparing each wetland's proportion "full" to percent of average total precipitation for time steps of the past 2, 4, and 8 years. A wetland's best correlated time step is reported for all $p < 0.10$.

3.2 CLIMATE

3.2.1 2009-2010 weather

I analyzed daily precipitation records for May through August of 2009 and 2010. Climate data are available since 1931 for the Mammoth and Tower weather stations (NCDC, 2011). All study wetlands are located within 12 km of one station and 350 m elevation of both stations

(Fig. 1). In 2010 a HOBO tipping bucket rain gauge (Onset Computer Corp.) was installed (UTM Zone 12N 549867 E, 4973438 N) to measure precipitation near Trumpeter Lake. I performed linear regression analysis on 2010 weekly precipitation data among the Tower, Mammoth, and HOBO rain gauges to analyze spatial variability among study sites.

3.2.2 Historic datasets

The Yellowstone River above the Corwin Springs gauging station (USGS gauge #06191500) drains 6783 km², including the entire Northern Range within YNP. Mean daily discharge data are available for the period 1911-2010. The shape-magnitude framework described above was used to analyze Yellowstone River discharge trends. To classify magnitude from each year, the maximum, minimum, mean, and standard deviation of monthly discharge data were used for annual hydrographs. The Palmer Drought Severity Index (PDSI), a metric of dryness (Dingman, 2002), is also reported for the Yellowstone River drainage basin from 1895-2010 (NCDC, 2011).

3.3 VEGETATION

3.3.1 Plant-hydrology relationship

Vegetation in the study basins was typically distributed in concentric zones controlled by water depth and duration. In 2009 I estimated the canopy cover of each plant species present in a 2 m x 50 cm plot centered on each well and oriented parallel to vegetation zones. To determine the hydrologic conditions supporting common plant species, I calculated the mean weekly water table depth for all species comprising $\geq 20\%$ cover at four or more wells. This included *Carex aquatilis*, *C. atherodes*, *C. pellita*, *C. utriculata*, *Eleocharis palustris*, *Phleum pretense*, *Poa pratensis*, *Schoenoplectus acutus*, and four *Salix* species, *S. boothii*,

S. drummondiana, *S. geyeriana*, and *S. pseudomonticola*. The *Salix* were all found in peatlands, had similar hydrologic niches, and were combined for analysis.

3.3.2 Soil seed bank study

Soil seed banks were investigated to assess a wetland's ability to revegetate if higher water levels return. Although some wetland plants reproduce asexually through rhizomes, many species produce persistent seed banks that stay viable for decades. The soil seed bank was analyzed at 12 wetlands that represented a variety of Northern Range environments. Soil samples were collected along two transects beginning at either the middle of the wetland basin or approximately 40 cm below surface water in deep wetlands. Transects extended to the surrounding upland vegetation. Each transect included five to seven plots spaced at 30 to 60 cm elevation intervals, corresponding to the vertical relief between wetland bottom and upland. A 25 x 25 x 5 cm thick soil block was collected from each plot at the ground surface, air dried, and sieved to remove roots and rhizomes. Each sample was spread onto sterile Pro-Mix BX soil in a Colorado State University greenhouse and analyzed using the seedling emergence method (Roberts, 1981). Soils were subjected to waterlogged conditions, which have been shown to be superior to ponding for eliciting germination of wetland species (Boedeltje et al., 2002). Vega and Sierra (1970) found that 83% of seedlings emerging in three years germinated in the first year. For this study all plants identifiable within 14 months were included.

For statistical analysis, plots on each transect were assigned a value from 0 to 1, representing their proportional elevation from the wetland's interior to the surrounding upland. This method produced a relatively even distribution of a given wetland's plots among elevation classes, and it preserved the role of scale in variously-sized watersheds. All 140 plots were

stratified into seven elevation classes that were analyzed using Kruskal-Wallis and Mann-Whitney U tests with the program SAS (2010).

3.4 TRUMPETER LAKE FOCAL SITE

3.4.1 Surveying and aerial photograph analysis

Because of its ecological significance and importance to the public, Trumpeter Lake was chosen as a focal site. A lichen trimline on glacial erratics around the lake indicates that lake stage was higher in the recent past (Marsh and Timoney, 2005; Hale, 1974). In northern Alberta, Marsh and Timoney (2005) showed that trimlines can persist 33-65 years. The declined lake water level has triggered soil erosion and habitat alteration detrimental to trumpeter swans and other native species. I used a total station to survey the zone between the water's edge and the maximum possible Trumpeter Lake stage. An existing bathymetric dataset was used to estimate land surface's elevation under surface water (Jones et al., 1978). Data were imported to ArcMap 10.0 where a triangulated irregular network (TIN) surface was created. Surface water polygons were derived from each of the eight air photos and superimposed on the TIN to determine lake stage in each photo. The topographic survey allowed me to analyze lake volume in addition to area (Hayashi and van der Kamp, 2000). To minimize seasonality bias, I analyzed lake sizes on a common day of year by calculating changes in lake stage between the photo date and August 15. To do this I subtracted the measured precipitation from evaporative loss, which was estimated using historic monthly (Pochop et al., 1985) and annual (WCA, 2008) pan evaporation measurements. This value was added to lake stage if the photo was taken before August 15, or subtracted from photo stage if taken after August 15. The calculated stage superimposed on the TIN produced lake sizes for a common day of the year. 2010 lake size was measured in the field.

To investigate the time scale on which Trumpeter Lake functions, regression analysis was used to compare lake area and volume to precipitation totals over the previous 1, 2, ..., 10 years.

3.4.2 Aquifer flow paths

The time lag between precipitation input and subsequent water level changes in the wetland basin is influenced by aquifer saturated hydraulic conductivity (K_s). Two methods were used to calculate K_s , the Hvorslev slug test (Fetter, 1994) measured water's return rate in groundwater monitoring wells in the Trumpeter Lake watershed. A double-ring infiltrometer (ASTM, 2003) was used to calculate K_s at the ground surface and at 50 cm depth on lake-margin and upland till environments. Water samples taken in the Trumpeter Lake watershed were analyzed for $\text{Ca}^{2+}/\text{Na}^+$ ratios. Differences in this ratio reflect different flow paths.

4. RESULTS

4.1 WETLAND HYDROLOGIC REGIMES

4.1.1 Wetland hydrology

Northern Range wetlands exhibited several indicators of surface water decline. At three sites, OR, LT, and TL, eroded former shorelines occurred > 200 cm in elevation above the 2009-2010 maximum surface water level (see Table 1 for wetland acronyms). Lichen trimlines, created when surface water drowned established lichen colonies (Marsh and Timoney, 2005; Hale, 1974), occurred on rocks surrounding wetland basins at BM (95 cm above maximum measured water), CR (105 cm), LT (230 cm), and TL (250 cm). These rocks had little or no visible lichen colonization below the trimline, suggesting that the high water period occurred within the past few decades (Timoney and Marsh, 2004). A third indication of former water depth was the location of dead stands of bulrush, an emergent marsh species. These slowly decaying plants

occurred on highly organic soils 20-30 cm thick and were 180, 185, and 200 cm above the study period's highest surface water at OR, LT, and TL.

During my study period surface water levels declined through the growing season at 22 of the 24 study wetlands, and all of these sites had lower water levels during 2010 than 2009 (Fig. 4). In 2009 all 24 study sites supported surface water in early summer and 15 retained it through mid-August, while in 2010, 22 sites had surface water in early summer and 9 did in mid-August. Water table variance during the summer differed by site. Wetlands DW and RL had persistent inflow and outflow, and surface water levels varied < 3 cm during both summers. In contrast, surface water level in BD, BG, and CP averaged 111-119 cm higher in 2009 than 2010.

Groundwater level patterns resembled those of surface water levels for most wetlands (Fig. 4d). Each summer's most intense two-week rain period, 29 mm in 2009 and 52 mm in 2010, triggered groundwater rises (Fig. 3, Fig. 4c). In some sites groundwater levels varied more than surface water (e.g. TL, BW), while at other sites the opposite occurred (e.g. BD, SV). 19 wetlands most frequently experienced groundwater inflow from one side and outflow to the opposite side, and thus were considered flow-through systems (Dingman, 2002). Hydraulic gradient reversals, where the relative elevations of groundwater and surface water reverse, were documented at seven sites (Fig. 5e). Following the periods of heaviest rain (e.g., the first two weeks of June 2010), LT and other wetlands switched from flow-through wetlands to having groundwater inflow on all sides (Fig. 4a). Wetlands MC, BP, and WA were perched above the regional water table.

Piezometric head data revealed that vertical gradients were small and groundwater flows were predominantly horizontal. Only 17 of 217 piezometers had significant upward gradients, characterized by a hydraulic head ≥ 5 cm from water table elevation. A spring supporting DW had 22-23 cm of positive gradient, and one supporting TF and TL had 25-48 cm

of positive gradient. 12 of 17 piezometers with positive gradients took longer than one week (median 10 weeks) to equilibrate following installation, suggesting they had low K_s and produced little upward flow into the wetland. Seven piezometers consistently had more than 5 cm of negative gradient, indicating that downward flow is uncommon. The three perched sites all had at least one deep piezometer that never contained water even when surface water was present, revealing an unsaturated zone below surface water.

Wetlands containing surface water outflow at any time during the study period were distinct from closed basins. These wetlands typically had inflowing groundwater, and their surface outlets maintained relatively stable maximum water levels. They had similar early summer water table levels in both study years. Five of the seven wetlands with outlets had peat soils and are fens (Lemly and Cooper, 2011), and the other two are a large and deep lake (RL) and a perched site (MC) that was often dry. Wetlands with outlets had less water table response to early summer rains (mean 2 cm) than closed-basins (13 cm; $t = 2.87$, $p = 0.005$, Fig. 4c&d). Sites with outlets responded to rain events in late summer only if surface outflow had ceased (Fig. 4d).

The Trumpeter Lake wetland is a flow-through hydrologic system, with groundwater inputs from the south and southeast (Fig. 5, Fig. 6). $\text{Na}^+/\text{Ca}^{2+}$ ratios from groundwater samples indicated that two separate aquifers contribute to water tables in the watershed. A spring from the bedrock aquifer had $\text{Na}^+/\text{Ca}^{2+}$ ratios around 13, while ratios from the surficial till aquifer springs were around three. Trumpeter Lake stage declined by 23 cm in 2009 and 45 cm in 2010. Seasonal water table maxima occurred in early June in both years. By late summer 2010, surface water was higher than groundwater on all sides of the lake and recharged groundwater on all fronts (Fig. 6d). Double-ring infiltrometer and slug tests in the Trumpeter Lake watershed indicated that K_s values ranged from 0.003 to 1.7 m/day (Appendix D). The double-ring

infiltrometer yielded K_s values roughly an order of magnitude higher than the slug tests, commensurate to the findings in till by van der Kamp and Hayashi (2009). These low conductivity values indicate that it can take years between water's arrival in the watershed and its interaction with the lake.

4.1.2 Wetland Classification

Seven wetland classes were identified from water table variation patterns. Description of these classes follows:

Stable surface water, "stable": DW, RL. Basins maintaining surface water within 3 cm of constant due to consistent inflow and outflow. Dense emergent vegetation bordered the surface water. Aerial photograph analysis indicated that wetland area has remained stable for decades.

Perched water tables, "perched": BP, MC, WA. Wetlands with perched water. Water levels steadily declined before disappearing. Some piezometers never contained water even though a nested monitoring well did, indicating the piezometer was completed in or below a confining layer.

Seasonally and yearly variable water tables, "seasonal": BD, BG, BM, BW, CP, CR, SV. Wetlands with large water table decline during and between summers. 2010 water levels averaged 75 cm lower than in 2009, more than double the between-year decline of any other class. Winter snowpack created flow-through or recharge basins, and pronounced water table rises following rain events indicated a close connection between precipitation and water tables.

Peat soils with recovering water tables, "P.R.": DC, RP, SN. Fens supported by groundwater discharge with water levels declining in late summer and recovering during the winter.

Recovering is defined by water tables increasing between study summers due to recharge processes. 2009 water level declines occurred later in the summer and were smaller than in 2010. All sites had surface water outlets that limited the water table's response to early summer rain.

Mineral soils with recovering water tables, "M.R.": MB, US. These mineral soil basin wetlands had the largest water table declines, approximately 100 cm each summer. Water tables recovered during the 2009-2010 winter.

Interannually variable water table persistence, "I.V. ": BE, IP, OR, SG, TF. Water tables declined minimally in 2009 but much more in 2010. Sites included those with and without outlets and having mineral and organic soils. Water table variation was relatively small and unresponsive to precipitation events.

Spatially variable water table patterns, "S.V.": TL, LT. Wetlands containing wells from at least three of the well types described above. Water tables in different areas of these wetlands varied distinctly, indicating that a combination of hydrologic patterns converge and create spatial complexity.

Wetland class assignment was influenced by the interaction between each wetland and its surrounding environment. Wetland class was correlated with the highly linked variables of surface water outflow ($\chi^2 = 16.03$, $p = 0.01$) and peat soil ($\chi^2 = 15.77$, $p = 0.02$), variables that were indicative of wetlands in classes P.R. and I.V. Surface water outlets constrained maximum wetland water levels, dampened water table changes, and supported peat formation. The most parsimonious classification tree in CART analysis contained only one split, divided at peat presence or absence, a variable 92.5% correlated with surface water outflow. The overall CART misclassification rate was 12.5%, compared to the majority misclassification rate of 29.2%.

Therefore, the model reduced misclassification by 16.7%. Basin size differed among wetland classes ($F = 3.5$, $p = 0.02$) and appeared to be driven by the largest two wetlands being both members of the S.V. class. Presence of a clay layer beneath the wetland basin ($\chi^2 = 14.1$, $p = 0.03$, Table 2) was most prominent in the perched class.

4.1.3 Wetland size through time

Aerial photographs from eight years for the period 1954 to 2009 indicated that surface water extent has varied up to 400% at study sites. Wetland areas were greatest in 1969 and 1998. 1969 followed the wettest decade preceding any air photo, and 1998 followed the two highest runoff years of record for the Yellowstone River. Wetlands with inlets and outlets (DW and RL) maintained constant area during the photo study period. Other sites varied substantially, with 10 of 14 wetlands completely lacking surface water in some years.

Surface area at BD, IP and CR was correlated with total precipitation over the two years prior to the photo date, while LT, TL, BM, and CP were correlated with precipitation over the previous eight years (Table 3). The remaining seven wetlands were not clearly correlated with precipitation patterns. Wetland surface area was more strongly correlated with total annual precipitation than with snowfall alone ($t = 2.70$, $p = 0.007$).

Trumpeter Lake's August 15 area peaked at 12.8 ha in 1969 and declined to 3.0 and 3.5 ha in 2006 and 2010 (Appendix E). A topographic survey revealed that Trumpeter Lake's volume was highly correlated with its area by a power function ($R^2 = 0.999$, Fig. 7). Lake volume was more highly correlated with precipitation over the previous 5-10 years than the previous 1-4 years alone (Table 4), suggesting that groundwater accumulated over several years sustains the lake.

For the 16 wetlands with an identified relict hydric soil, this soil's upper boundary differed significantly among wetland classes compared to the current surface water level ($F = 3.29$, $P = 0.04$). This finding provides a longer-term perspective supporting the wetland classification, which was developed from two summers of hydrologic data. Hydric soil boundaries at wetlands with an outlet were more similar to the 2009-10 water levels ($n = 7$, mean = $31 \text{ cm} \pm 4 \text{ cm SE}$ above 2009-10 mean peak) compared to closed-basin sites ($n = 9$, mean = $128 \pm 24 \text{ cm SE}$, Student's t-test, $t = 3.31$, $p = 0.005$). The presence of hydric soils in locations beyond the surface water elevations measured during field study provided evidence suggesting that wetlands occupied greater areas in the past.

4.2 CLIMATE

4.2.1 Trends in Yellowstone River discharge

The 5-year mean discharge of the Yellowstone River was positively correlated with wetland area over time ($R = 0.89$; Fig. 8a), suggesting that the river's discharge record from 1911-2010 may be a good proxy for wetland area. Cluster analyses of annual Yellowstone River discharge data produced three magnitude and four shape classes, the latter corresponding to annual runoff peaks occurring in May, June-May, June, and June-July (Fig. 9). In the 26 years since 1984 there have been no late peak and 10 early peak years, compared to their relatively equal distribution of runoff timing in the decades prior to the 1980s (Fig. 10). The ratio of May (early) to June-July (late) peaks in the last three decades is unprecedented in the discharge record. Prior to 1984 the mean annual discharge peak occurred on June 11, but from 1985-2010 it occurred on May 30 (t-test, $t = 4.67$, $p < 0.001$). The decade most similar to the early peak-dominated 1990s and 2000s was the drought period of the 1930s. This decade was followed by the 1940s, which had the most late peak years on record. The early peak 1990s, however, was

followed by a second successive early peak decade. Unlike discharge timing, the volume of mean annual discharge did not change pre- and post-1984 (Student's t-test, $t = 0.38$, $p = 0.70$).

4.2.2 Long-term climate trends

For the last 100 years, relative highs and lows in Yellowstone River discharge and PDSI have co-occurred (Fig. 8). Since 1970 river discharge has oscillated around the mean of the past century, while increased temperatures have produced consistently lower PDSI values. Although PDSI trends suggest that the past four decades have been drier than most of the last century, tree ring data indicate that conditions at least as dry as the past 40 years have occurred in YNP several times in the past 800 years, for example *circa* 1250, 1500, 1700, and 1800 AD (Fig. 8d; Gray et al., 2007). The relatively dry second half of the 20th century was preceded by the most prolonged wet period of the past 800 years, lasting approximately 80 years from the late 19th through the early 20th centuries (Gray et al., 2007). This wet period may have resulted in the expansion of Northern Range wetlands to their largest extent in the past several centuries.

4.3 VEGETATION

4.3.1 Vegetation-hydrology relationship

The nine most common plant species in study wetlands occupied distinct hydrologic regimes (Fig. 11). *Carex aquatilis* and *C. utriculata* occurred in locations with the highest water tables in both years, having surface water throughout the first half of each summer. *Carex pellita*, *Schoenoplectus acutus*, and *Salix* spp. were also found in wet locations with water tables near the soil surface through mid-June in 2009. In 2010 their water tables were 10-20 cm below ground in early summer and dropped to approximately 75 cm in late summer. *Eleocharis palustris* occurred in locations that were saturated in early 2009, but the water tables dropped

substantially in both years. *Carex atherodes* experienced similarly large summer variation as *E. palustris*. It occupied sites with mean 2009 and 2010 water tables differing by 60 cm. *Phleum pratense* and *Poa pratensis* occupied the driest locations. The mean water level at sites supporting *Poa pratensis* was 42 cm lower in 2010 compared to 2009, while *Phleum pratense*'s was 23 cm lower. *Poa pratensis* inhabited zones of fluctuating hydrologic conditions, while *Phleum pratense* inhabited more consistently dry sites.

Carex-dominated wetlands, excluding fens, had outlets within 1 m of measured surface water maxima. In contrast, bulrush-dominated sites lacked outlets within 1 m (*non-significant*, $p = 0.12$). During large water periods bulrush sites can support much deeper water levels than those dominated by *Carex*. Bulrush sites had less water table difference between study years indicating that wetlands supporting this species may have greater year-to-year stability. Aerial photos, hydric soils, and outlet elevations all suggest that bulrush tolerates larger decadal water level changes compared to *Carex*. *Carex*-wetlands were often members of the seasonal wetland class.

This study identified two wetland plants not previously recorded in YNP. *Scirpus nevadensis* S. Watson was found at wetlands LT and IP, and *Alopecurus geniculatus* L. was found at WA.

4.3.2 Soil seed bank study

46 plant species germinated from soils collected at the wetland study sites. Six species were obligate wetland species (OBL) and nine were facultative wetland species (FACW; Appendix F). In a given plot, some species were represented by one germinant, while others had hundreds of germinants. The relative elevation-based classification produced a marginally significant unimodal curve for the distribution of OBL and FACW wetland species (Kruskal-Wallis

test, chi-square = 11.76, df = 6, $p = 0.07$). Significantly more wetland species germinated just above the midpoint of the wetland bottom and upland boundary compared to the lowest three and highest classes (Mann Whitney U test, $z = 2.00-2.88$, $p = 0.01-0.05$; Fig. 12).

5. DISCUSSION

Multiple lines of investigation helped identify the patterns and processes driving wetland losses on YNP's Northern Range. Wetland area and water level is clearly linked to weather and climate patterns, and the wetland sizes measured in this study have likely occurred multiple times during the past millennium. The first decades after YNP's designation as a national park coincide with the most prolonged wet period of the past 800 years, which may have produced unusually large wetlands. YNP's climate has also warmed and dried during the last century, and continued warming could surpass the natural range of variation that occurred within the tree ring record and could lead to unprecedentedly small wetland areas. Wetlands serve a valuable role in maintaining ecosystem integrity in YNP, which is one of the last places in the conterminous United States where natural ecosystem processes dominate. Declines in YNP's native species have been linked to wetland reduction (McMenamin et al., 2008; Proffitt et al., 2010) and continued drying will endanger additional species and cause further environmental changes, such as shifting fens from carbon sinks to carbon sources (Chimner and Cooper, 2003).

Each study wetland's local watershed controlled its hydrologic regime. This is distinct from the influence of the broader regional hydrologic patterns identified in other wetland settings (Florin et al., 1993; Merkey, 2006; Fig. 1). Wetland areas vary with climate, including those in YNP, the prairie potholes (Johnson et al. 2004, Niemuth et al. 2010), southern Alaska (Klein et al. 2005), and Siberia (Smith et al. 2005). Similar to wetlands in Argentina's Monte Desert (Jobbágy et al., 2011) and Nebraska's Sandhills (Winter, 1986), groundwater sustained

many YNP study wetlands. A few wetlands lacked groundwater contributions, similar to playas of the southern American Great Plains (Smith, 2003) and vernal pools in California (Keeley and Zedler, 1998). Two study wetlands have apparently maintained stable water levels for decades, while others have varied substantially. Rarely do wetlands in such close proximity function as differently as has been documented here. The various hydrologic regimes of Northern Range wetlands influence local ecosystem processes including soil microbial environments and lichen, vascular plant, and animal communities.

5.1 HYDROLOGICALLY-BASED WETLAND CLASSIFICATION

Classification was a practical tool for creating a conceptual framework to compare Northern Range wetland hydrologic regimes. Study wetlands differed to a degree approaching the national-scale variability documented in classifications from the United States (Brinson et al., 1993; Cowardin et al., 1979), Spain (Florin et al., 1993) and Amazonia (Junk et al., 2011). The shape-magnitude framework incorporated the important roles of timing and degree of water table changes to identify seven wetland hydrologic classes. Hydric soils analysis substantiated the wetland classification by showing that the identified classes have functioned distinctly for decades. Wetlands from the seasonal class provide habitat for species adapted to changing surface water abundance, such as *Carex atherodes*, *Carex utriculata* and YNP's four native amphibians. The S.V. wetland class included large wetlands with multiple water sources. Consistent saturation has allowed P.R. wetlands to accumulate organic matter and create peat soils. Three of the five wetlands in the I.V. class were peatlands with groundwater inputs that persisted for longer than those in the P.R. class. The two other I.V. sites were both closed basins with mineral soil, but they differed from the M.R. and seasonal classes by having more persistent groundwater inputs. Wetlands in the stable class were uncommon across the

Northern Range but may serve as isolated habitat refuges for wetland-dependent flora and fauna during exceptionally dry times.

5.2 HYDROLOGIC PROCESSES CONTROLLING YELLOWSTONE'S WETLANDS

Inputs to wetland water budgets consisted of surface water, groundwater, and precipitation. Rain produced a relatively rapid response in wetland sizes, while snow slowly recharged aquifers and wetted soils to initiate the growing season. The groundwater discharge supporting 21 of 24 study sites stabilized water tables, presumably producing longer growing seasons and increasing vegetative production (Rains, 2011; Cook and Hauer, 2007; Hunt et al., 1999). Groundwater flow paths were not fully characterized because land surface watersheds can differ from groundwater watersheds (Devito et al., 2005) and multiple wetlands shared nested watersheds in the study area's poorly developed land surface drainages.

Watershed characteristics including topography, lithology, and infiltration capacity strongly influence hydrologic processes in wetlands (Florin et al., 1993) and vary on small scales in the Northern Range. Peat soil and the presence of a surface water outlet were the two environmental variables most closely correlated with wetland class in this study. Surface outlets limited maximum water level and increased hydrological and ecological stability. The resulting near-surface water table promoted high plant production, low decomposition rates, and peat formation. Merkey (2006) incorporated outlet presence into a wetland classification for Michigan, finding that sites with both inlets and outlets exhibited unchanging water tables, as occurred in my stable wetland class. At study wetlands DC and SG, surface water inflow and outflow existed seasonally, but limited inputs caused water tables to decline by late summer.

5.3 HYDROLOGIC CONTROLS OF WETLAND VEGETATION

The distribution of common wetland plants indicated both past and present hydrologic regimes. Robust bulrush stands were found only where relatively stable standing water levels occurred in both study years. Decomposing dead bulrush stands with thick organic soil occur at three sites > 2 m above the highest measured surface water level. Two of these sites are classified into the S.V. wetland type and responded to the long-term patterns of precipitation. Bulrush appears to be a good indicator of wetland basins whose water levels change slowly, but that can change substantially in different climate periods.

Dominant wetland plant species occupied distinct zones of water depth and seasonal variation. Because each species occupied a distinct hydrologic niche, point measurements from monitoring wells in patches dominated by common species can be used to extrapolate hydrologic processes to other areas where the species are found. Plant community structure differs in areas of groundwater discharge and recharge (Cook and Hauer, 2007; Hunt et al., 1999; Humphries et al., 2011) and can indicate the spatial distribution of these hydrologic processes at a wetland. Wetland vegetation will change as the climate changes (Johnson et al., 2010; Poiani et al., 1996), therefore monitoring plant distribution at study sites may help identify the ecological effects of climate change. Plants germinating from the soil seed bank indicated that when higher water levels occur in the future, viable seed banks can promote rapid wetland vegetation adjustment.

5.4 TRUMPETER LAKE WATERSHED DYNAMICS

Lichen trimlines, relict bulrush stands, and an eroded shoreline reveal a 2.5 m surface water decline at Trumpeter Lake during the past few decades. Sediment cores have been used to show that water levels are highly variable and suggest that the lake has been dry at some point in the last century (Engstrom et al., 1991). This water table change parallels the “drought

and deluge” conditions in semi-permanent wetlands of the prairie potholes (Johnson et al., 2004). K_s values in the Trumpeter Lake watershed were on the order of dm/d at the land surface and cm/d at 1-2 m depth, similar to values recorded in glacial till in the United Kingdom (Cuthbert et al., 2010) and the prairie pothole region (van der Kamp and Hayashi, 2009). Higher K_s surface values may result from the abundance of near-surface macropores created by root channels and burrowing animals (Cuthbert et al., 2010; Todd et al., 2006). Near-surface K_s values encompass only a fraction of the till’s thickness (K. Pierce, *pers. comm.*), but the bulk of water movement is likely within this section due to decreasing K_s with depth in till aquifers (van der Kamp and Hayashi, 2009). Groundwater discharging southeast of Trumpeter Lake may be emerging where a zone of thin till permits water expulsion from the underlying bedrock (Todd et al., 2006). However, chemical analysis of this water indicated that its origin is more likely from the surficial till aquifer. A model of the lake’s water budget would help to analyze lake volume variations, but a robust model calls for greater resolution of aquifer thickness, K_s variability, and watershed ET.

The highest recorded population of Northern Range bison (Wallen, 2010) appears to be having destructive effects on the Trumpeter Lake area vegetation and soil, likely driving vegetation changes, denuding and eroding the land surface, and altering water infiltration. Bison hooves sink > 30 cm into lake edge soil, promoting erosion. Rocks located between the former and current lake perimeters have rust-colored stains on their bases, suggesting they were recently beneath the soil surface and have been exposed by erosion. In addition, dead bulrush 2.0 m above the modern lake surface occurs in isolated clumps 20 cm above the surrounding ground surface. All contemporary bulrush stands are comprised of homogenous clones that cover the ground in dense continuous carpets, not clumps, indicating that a strong root system and slow decomposition have prevented soil erosion at the dead bulrush locations. Additionally,

vegetation colonizing the zone between former and current Trumpeter Lake perimeters consists of low-diversity stands of invasive *Hordeum jubatum*, *Cirsium arvense*, and *Argentina anserina*.

5.5 WEATHER, CLIMATE, AND WETLAND AREA CHANGES THROUGH TIME

Evidence on the Northern Range indicates that wetland water levels have declined, but the processes of drying have varied by wetland. Seasonal wetlands had the greatest stage changes during 2009-2010 and responded to early summer rains with >50 cm water table increases. In contrast, wetlands in the stable, P.R., and I.V. classes exhibited little response to rain events. These sites either had surface water outlets or were closed basins that buffered response to individual precipitation inputs. Aerial photograph analysis indicated substantial variability in water abundance, suggesting that in the future air photos should be taken over broad spatiotemporal scales and at the same time each year (Niemuth et al., 2010). Some wetlands revealed undetectable connections to precipitation in aerial photograph analysis, even though it undoubtedly affects them. This could be due to the time scale on which a wetland's hydrologic regime functions being outside of the scale analyzed in aerial photos, or due to interacting landscape factors. For example, WA is a basin located at the base of a bedrock outcrop that may accumulate wind-blown snow (e.g., Rains, 2011).

Although watershed properties play an integral role in creating a wetland's hydrologic regime, it is likely that at larger time scales climate is more important (Laize and Hannah, 2010). Data from the past century support the idea that climate change is responsible for wetland loss. A warming trend has occurred in YNP, including exacerbated changes since the 1980s documented here and by others (McMenamin et al., 2008; Wilmers and Getz, 2005). Johnson et al. (2010) modeled climate change's effects on wetlands of variable permanence, finding that the less permanent sites (e.g. seasonal) are more resilient to climate change than more

permanent ones (e.g., S.V. and P.R.). This was because of the latter's support from groundwater and increased evapotranspiration losses from surface water persisting through summer. Seasonal class wetlands can change from their maximum water extent to dry within 1-2 years, but their flora and fauna are adapted to this variability. However, it is likely that during the recent multi-decadal drying, P.R., S.V., and I.V. class wetlands have become drier as well. Under projected climate changes, the growing season's start and the timing of seasonal water table decline will occur earlier, leading to additional drying, organic matter decomposition, and wetland destabilization. Non-linear or threshold responses (Burkett et al. 2005) may limit fens' ability to sequester carbon. Climate data reveal that during the past 115 years YNP has warmed and dried, but tree ring data from Gray et al. (2007) through 1998 show that the recent dry spell has been surpassed multiple times in the past 800 years. Since 1998 the Northern Range has recorded a mean of 86% of historic annual precipitation, revealing further drying since this dataset was collected. However, it is unlikely that 12 moderately dry years have shifted the system to an unprecedented state. Gray et al. (2007) also document that the late 19th and early 20th centuries experienced the most prolonged wet cycle of the last 800 years. This suggests that efforts to revert Northern Range wetland abundance to the pre-climate change era may be a misguided attempt to reclaim an exceptionally wet "reference state".

Wetland changes have been poorly documented because of the multitude of wetlands on the landscape and the investment required to monitor even a single site. Lentic wetlands and lotic waterways, although inherently different, each are driven by groundwater regimes that are influenced by climate. This study suggests that Yellowstone River discharge data is a useful proxy for wetland surface water area on the Northern Range. Comparison in other regions should investigate the generality of this relationship. This study and others (McMenamin et al., 2008; Stewart et al., 2005) have shown Rocky Mountain river hydrographs shifting to earlier

peaks, and Stewart et al. (2004) projected further advancement by 30-40 days throughout the West. This hydrologic shift would initiate major ecosystem changes in wetlands, where the water table position relative to the land surface is critically important to ecological processes.

6. CONCLUSIONS

Wetlands on Yellowstone's Northern Range have declined in area during the recent past, but the evidence suggests that current conditions are within the natural range of variation for the past 800 years. Furthermore, it is often assumed that when Westerners arrived, YNP's landscape was in a relatively pristine state of equilibrium. However, this was a period of water abundance. Weather patterns naturally oscillate between warm-dry and cool-wet conditions, and the warmer and drier conditions of the last several decades have reduced wetland area and surface water levels. If the recent climate trajectory continues, wetlands will dry further, removing critical habitat for native species.

The research approach taken here has created a multifaceted perspective of YNP's recently dried wetlands. A combination of hydrology, soils, vegetation, and climate investigations were used to synthesize why, how, and to what degree wetland changes have occurred. Monitoring well data, aerial photographs, hydric soils, bulrush vegetation, and lichen trimlines all illuminated the patterns and magnitudes of wetland changes. The hydrologic wetland classification aided in creating an understanding of why these patterns have occurred. Wetlands in different classes are differentially vulnerable to forecasted climatic changes, and this should be considered in management. Wetlands from classes I.V., S.V., M.R., and P.R. were relatively resilient to the different 2009 and 2010 weather patterns, indicating that longer-term climatic processes more likely affect their water levels. If YNP climate continues to warm and

dry, wetlands in these classes and their associated ecological functions (e.g. carbon sequestration, amphibian breeding, trumpeter swan nesting, etc.) will be vulnerable.

Preservation of landscape-scale ecological processes is a primary goal of the National Park Service, and wetland health serves a central role in this endeavor. Two years of data were collected in this study, but a thorough hydrologic study necessitates longer-term data collection and analysis. My analyses indicate that wetland conditions such as those in the S.V. class may be relatively common; however, decadal changes have affected our perception of the significance of their drying. Wetlands from the seasonal, P.R., and I.V. classes formed a continuum from low to high permanence. If these wetlands shift down a permanence class it could have dramatic ecosystem consequences, including loss of peatlands that formed over millennia.

Climate change was the primary candidate causing wetland change addressed here, but other factors should also be considered. YNP's dynamic geology is a force underlying all surficial processes in the Northern Range, and unprecedentedly high bison populations affect wetlands by increasing the rate of soil erosion and vegetation change. Topsoil erosion should be monitored in the Little America area because losing this relatively fertile soil may further degrade these wetland ecosystems. Wetlands serve a critical role on the Northern Range landscape, and we should continue to monitor them to recognize if-and-when conditions surpass the ecosystem's capacity to adjust.

7. FIGURES AND TABLES




Figure 1. The Northern Range within Yellowstone National Park, Wyoming and Montana, USA.

Study sites are represented by dots and colored according to their water table hydrograph classification. Mammoth and Tower weather stations are represented by M and T.




Figure 2. Cluster analysis results for the 80 wells analyzed using the shape-magnitude framework. A) dendrogram of hydrograph shape with pruning at 50% similarity which produced 3 classes. B) dendrogram of magnitude with pruning at 40% similarity which produced two classes. Note that the first two letters in well names are wetland IDs.




Figure 3. Water table elevations for the seven well classes during summers 2009 and 2010.

Lines are average weekly water table values for all wells in each class, except for the perched class which depicts an example well that dried in June 2010. The S.V. wetland type described in the Results was comprised of at least three of these well types. Each of these well classes was developed into a unique wetland class except shape1-mag2, which was too uncommon (n = 6) to constitute its own class. Bars represent weekly precipitation.




Figure 4. Panes A & B show cross sections of wetlands LT and DC. Solid lines represent ground surface, and dashed lines are water table elevations on 10 June 2010 (grey) and 5 August 2010 (black). Water tables were interpolated from groundwater monitoring wells (thick bars). Piezometers (thin bars) nested with wells reveal vertical hydrologic gradients. Note that piezometers at left in pane A show an upward gradient, but all other piezometers show nominal vertical gradients. Bars indicate water level on 10 June (gray) and 5 August (black). Panes C and D are hydrographs showing water level in wells and piezometers from A and B during summers 2009 and 2010. Solid lines are wells, and dotted and dashed lines are piezometers nested with the well of the same color. Hydrograph colors correspond to the same colored wells illustrated as dots on plan view insets in panes A and B. Labels in C and D correspond to well types; wetland LT was a S.V. wetland and DC was a P.R. wetland. Hydrographs revealed characteristics of site hydrology, including locations that: spiked in response to rain (blue in C and D), declined earlier in the drier 2010 (pink in C, all in D), had strong positive head (blue in C), had slowly-equilibrating piezometers that revealed a low hydraulic conductivity layer (pink dash-dot in D), and did not spike in response to early summer rain because of surface outflow (all in D).




Figure 5. Cross section of Trumpeter Lake illustrating the water table on four dates in 2010.

Ground surface, the thick line, was determined from a topographic survey. Grey bars represent monitoring wells, identified with stars on plan view inset showing all wells, which were used to interpolate water tables. Trumpeter Lake had groundwater inflow from the south in early summer, but by late summer a near-shore groundwater trough appeared. This indicates that the lake shifted from a groundwater flow-through system to groundwater recharge on all sides.

Vertical exaggeration is 80:1.




Figure 6. Trumpeter Lake water table contours (m.a.s.l.) for summer 2010, interpolated from 15 monitoring wells. Through 15 July this was a flow-through wetland, with the water table gradient from south to north. D) shows that in late summer surface water locally recharged groundwater (black arrows), even though the larger flow system was still flow-through. Water table elevation declined as the summer progressed beyond the early-June peak. Multiple groundwater interpolation methods were explored (e.g. IDW, kriging, spline), however, I manually delineated water table contours as this method is frequently more accurate (Bill Sanford, *pers. comm.*).




Figure 7. Surface area-volume relationship at Trumpeter Lake during nine years. Areas were derived from aerial photography (n = 8) and field data (n = 1). Corresponding volumes were calculated by overlaying the lake area on a TIN.




Figure 8. Multiple climate indices for YNP. A) Bars depict the across-wetland average % of maximum wetland surface area seen on air photos from a given year. Solid line is the 5-year running mean for annual discharge of the Yellowstone River at Corwin Springs. B) 5-year mean Palmer Drought Severity Index (PDSI) for the Yellowstone River drainage, Wyoming. C) 20th century 5-year mean tree-ring reconstructed precipitation for the YNP region (data from Gray et al. 2007)). D) Tree-ring reconstructed precipitation since 1173 AD, displayed as a 60-yr cubic smoothing spline (reproduced from Gray et al. 2007). Horizontal grey lines represent means

during the plotted period in A, C, and D, while in B it depicts the PDSI average by definition, 0.
 Note the unique x-axis for D.




Figure 9. Yellowstone River discharge shape-magnitude analysis classes. A) Mean annual hydrograph of all wells constituting each of the four shape classes, termed according to their timing of maximum discharge. B) Mean annual hydrograph of the 3 magnitude classes, big, medium, and small.




Figure 10. Chronology of the earliest and latest Yellowstone River runoff years, showing that early runoff has been unusually prevalent recently. Note: 1) Bars do not sum to 10 because the two middle classes were omitted for clarity; 2) The decade most closely resembling the 1990s and 2000s is the 1930s, the Dust Bowl years; 3) There is substantial inter-decadal fluctuation and no apparent 100-year trend. However, the 1970’s forward reveals a “slingshot” from late to early peaks, presumably inciting wetland decline.




Figure 11. Water table hydrographs for the nine most prevalent plant species. Graphs illustrate species' water table means \pm 1 standard deviation. Data are derived from averaging water tables for all wells with \geq 20% cover of the identified species.




Figure 12. Mean number of wetland species \pm se germinating per plot in the soil seed bank study. Bars represent the seven elevation classes, with the x-axis stratifying plots by their percent from wetland bottom to upland. Mann Whitney U tests revealed that plots in the 60-67% class germinated significantly more ($p < 0.05$) wetland plants than plots in the lowest three and highest elevation classes.

Table 1. The 24 Northern Range study sites.

Wetland Name	Wetland Acronym	Elev. (m)	UTM E	UTM N	n wells	Wetland Class
Big D	BD	1875	546592	4974876	4	seasonal
Bighorn Marsh	BM	1904	547949	4973587	6	seasonal
Blue-Green	BG	1905	545970	4974936	5	seasonal
Brown Pond	BP	1765	520612	4985821	3	perched
Bunsen East	BE	2218	522447	4973863	5	I.V.
Bunsen West	BW	2228	522035	4974291	5	seasonal
Chorus Pond	CP	2241	520797	4971697	4	seasonal
Copper Rock	CR	1887	550268	4973075	5	seasonal
Dead Willow	DW	1776	520432	4985701	7	stable
Double Cub	DC	2009	543547	4976370	4	P.R.
Island Pond	IP	1890	552001	4974167	5	I.V.
Little Trumpeter	LT	1861	549423	4973951	5	S.V.
Mammoth Bowl	MB	1824	524980	4978500	4	M.R.
Meadowlark Commons	MC	1769	520777	4985721	5	perched
Old Road	OR	1791	523300	4983570	5	I.V.
Rainbow Lake	RL	1792	520423	4985457	3	stable
Rye Pond	RP	1887	524180	4978560	5	P.R.
Sandhill Nest	SN	1864	548510	4974510	5	P.R.
Self-Guiding	SG	2047	534864	4979149	5	I.V.
Slough View	SV	1888	552075	4974492	5	seasonal
Trumpeter Feeder	TF	1863	550155	4973559	4	I.V.
Trumpeter Lake	TL	1860	549898	4973760	14	S.V.
Upper Slide	US	1752	523565	4983271	4	M.R.
The Wallows	WA	1884	551342	4973520	6	perched

Table 2. Relationship between environmental variables and wetland hydrograph classes. Chi-squared analysis was performed on categorical variables, and one-way ANOVA produced an F-stat for quantitative variables. Both tests produced comparable p-values (J. zumBrunnen, *pers. comm.*)

Variable	Categ/Quant	Chi²	F-Stat	P
Surface water outflow	C	16.03		0.014
Thick peat	C	15.77		0.015
Basin size	Q		3.5	0.02
Clay in basin	C	14.1		0.03
Pz. time to equilib.	Q		2.1	0.11
Watershed Size	Q		1.92	0.14
Highest pz. head	Q		1.75	0.17
Organic soil in basin	C	8.72		0.19
Elevation	Q		1.45	0.25
Precipitation	Q		1.15	0.37
Surface water inflow	C	6.24		0.4
Bulrush dominant	C	5.49		0.48
Basin EC	Q		0.9	0.52
Inflowing EC	Q		0.84	0.56

Table 3. Relationship between wetland area and cumulative precipitation, determined from aerial photo analysis. Wetland area and precipitation from the 2, 4, and 8 years previous to photo date were compared via forward-step multiple regression for the eight photo years. Results for a wetland’s most significant time step are reported if $p < 0.10$. No wetlands were most closely related to the last 4 years.

Wetland	Time step (yrs)	F-value	P-value
BD	0-2	8.32	0.028
IP	0-2	6.36	0.045
CR	0-2	3.92	0.095
LT	0-8	19.28	0.005
TL	0-8	58.19	0.0003
BM	0-8	9.17	0.023
CP	0-8	6.73	0.041
BG	none	-	-
BP	none	-	-
BW	none	-	-
WA	none	-	-
OR	none	-	-
DW	none	-	-
RL	none	-	-

Table 4. Relationship between total precipitation and Trumpeter Lake’s area and volume for time steps of the last 1, 2, ... 10 years. Data were derived from lake sizes in nine years.

Ppt. time step	Area R^2	Vol. R^2
1 Yr	0.33	0.47
2 yr	0.57	0.65
3 yr	0.78	0.79
4 yr	0.79	0.77
5 yr	0.86	0.81
6 yr	0.9	0.82
7 yr	0.92	0.87
8 yr	0.95	0.94
9 yr	0.89	0.87
10 yr	0.81	0.77

8. REFERENCES

- Acreman, M.C., Blake, J.R., Booker, D.J., Harding, R.J., Reynard, N., Mountford, J.O., Stratford, C.J., 2009. A simple framework for evaluating regional wetland ecohydrological response to climate change with case studies from Great Britain. *Ecohydrology* 2, 1-17.
- ASTM International, 2003. D3385-03: Standard test method for infiltration rate of soils in field using double-ring infiltrometer, 7 p., Washington, D.C.
- Bartlein, P.J., Whitlock, C., Shafer, S.L., 1997. Future climate in the Yellowstone National Park region and its potential impact on vegetation. *Conservation Biology* 11 (3), 782–792.
- Bates, B.C., Kundzewicz, Z.W., Wu, S., Palutikof, J.P., 2008. *Climate Change and Water*. Technical Paper of the Intergovernmental Panel on Climate Change, IPCC Secretariat, Geneva, 210 p.
- Belcher, W.R., Bedinger, M.S., Back, J.T., Sweetkind, D.S., 2009. Interbasin flow in the Great Basin with special reference to the southern Funeral Mountains and the source of Furnace Creek springs, Death Valley, California, US. *Journal of Hydrology* 369 (1-2), 30-43.
- Boedeltje, G., ter Heerdt, G., Bakker, J., 2002. Applying the seedling-emergence method under waterlogged conditions to detect the seed bank of aquatic plants in submerged sediments. *Aquatic Botany* 72 (2), 121-128.
- Bower, D., Hannah, D.M., McGregor, G.R., 2004. Techniques for assessing the climatic sensitivity of river flow regimes. *Hydrological Processes* 18, 2515-2543.
- Brinson, M.M., 1993. A hydrogeomorphic classification for wetlands. U.S. Army Corps of Engineers Wetlands Research Program Technical Report WRP-DE-4. 103 p.
- Brooks, R.T., 2000. Annual and seasonal variation and the effects of hydroperiod on benthic macroinvertebrates of seasonal forest (“vernal”) ponds in central Massachusetts, USA. *Wetlands* 20, 707-715.
- Brooks, R.T., 2004. Weather-related effects on woodland vernal pool hydrology and hydroperiod. *Wetlands* 24 (1), 104-114.
- Brooks, R.T., 2009. Potential impacts of global climate change on the hydrology and ecology of ephemeral freshwater systems of the forests of the northeastern United States. *Climatic Change* 95, 469-483.
- Carpenter, S.R., Fisher, S.G., Grimm, N.B., Kitchell, J.F., 1992. Global change and freshwater ecosystems. *Annual Review of Ecology and Systematics* 23, 119–139.
- Chimner, R.A., Cooper, D.J., 2003. Influence of water table levels on CO₂ emissions in a Colorado subalpine fen: an in situ microcosm study. *Soil Biology and Biochemistry* 35 (3), 345-351.

- Cook, B.J., Hauer, F.R., 2007. Effects of hydrologic connectivity on water chemistry, soils, and vegetation structure and function in an intermontane depressional wetland landscape. *Wetlands* 27 (3), 719–738.
- Cowardin, L.M., Carter, V., Golet, F.C., LaRoe, E.T., 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Department of the Interior, Fish and Wildlife Service, Washington, D.C. 131 p.
- Cuthbert, M.O., Mackay, R., Tellam, J.H., Thatcher, K.E., 2010. Combining unsaturated and saturated hydraulic observations to understand and estimate groundwater recharge through glacial till. *Journal of Hydrology* 391 (3-4), 263-276.
- Dahl, T.E., 2011. Status and trends of wetlands in the conterminous United States 2004 to 2009. U.S. Department of the Interior, Fish and Wildlife Service, Washington, D.C. 108 p.
- De'ath, G., Fabricius, K., 2000. Classification and regression trees: A powerful yet simple technique for ecological data analysis. *Ecology* 81 (11), 3178-3192.
- Devito, K., Creed, I., Gan, T., Mendoza, C., Petrone, R., Silins, U., Smerdon, B., 2005. A framework for broad-scale classification of hydrologic response units on the Boreal Plain: is topography the last thing to consider? *Hydrological Processes* 19 (8), 1705-1714.
- Dingman, S.L., 2002, *Physical Hydrology*. Vaveland Press, Long Grove, IL.
- Engstrom, D.R., Whitlock, C., Fritz, S.C., Wright, H.E., 1991. Recent environmental changes inferred from the sediments of small lakes in Yellowstone's northern range. *Journal of Paleolimnology* 5 (2), 139-174.
- ESRI, 2010. ArcGIS Desktop. Redlands, CA.
- Fetter, C.W., 1994. *Applied Hydrogeology*. Macmillan College, New York.
- Florin, M., Montes, C., Rueda, F., 1993. Origin, hydrologic functioning, and morphometric characteristics of small, shallow, semiarid lakes (lagunas) in La Mancha, central Spain. *Wetlands* 13 (4), 247-259.
- Gardner, P.W., Susong, D.D., Solomon, K.D., Heasler, H., 2010. Snowmelt hydrograph interpretation: Revealing watershed scale hydrologic characteristics of the Yellowstone volcanic plateau. *Journal of Hydrology* 383 (3-4), 209–222.
- Gee, G., Bauder, J., 1979. Particle-size analysis by hydrometer - simplified method for routine textural analysis and a sensitivity test of measurement parameters. *Soil Science Society of America Journal* 43 (5), 1004-1007.
- Gray, S.T., Graumlich, L.J., Betancourt, J.L., 2007. Annual precipitation in the Yellowstone National Park region since AD 1173. *Quaternary Research* 68 (1), 18–27.
- Hale, M.E., 1974. *The Biology of Lichens*. Edward Arnold, London.

- Hannah, D., Smith, B., Gurnell, A., McGregor, G., 2000. An approach to hydrograph classification. *Hydrological Processes* 14 (2), 317-338.
- Hayashi, M., van der Kamp, G., 2000. Simple equations to represent the volume-area-depth relations of shallow wetlands in small topographic depressions. *Journal of Hydrology* 237 (1-2), 74-85.
- Humphries, M.S., Kindness, A., Ellery, W.N., Hughes, J.C., Bond, J.K., Barnes, K.B., 2011, Vegetation influences on groundwater salinity and chemical heterogeneity in a freshwater, recharge floodplain wetland, South Africa. *Journal of Hydrology* 411 (1-2), 130-139.
- Hunt, R., Krabbenhoft, D., Anderson, MP, 1996. Groundwater inflow measurements in wetland systems. *Water Resources Research* 32 (3), 495-507.
- Hunt, R., Walker, J., Krabbenhoft, D., 1999. Characterizing hydrology and the importance of ground-water discharge in natural and constructed wetlands. *Wetlands* 19 (2), 458-472.
- Iglesias, P., Jorquera, H., Palma, W., 2006. Data analysis using regression models with missing observations and long-memory: an application study. *Computational Statistics & Data Analysis* 50 (8), 2028-2043.
- Jobbágy, E., Noretto, M., Villagra, P., Jackson, R., 2011. Water subsidies from mountains to deserts: their role in sustaining groundwater-fed oases in a sandy landscape. *Ecological Applications* 21 (3), 678–694.
- Johnson, W. C., Werner, B., Guntenspergen, G.R., Voldseth, R.A., Millett, B., Naugle, D.E., Tulbure, M., Carroll, R.W.H., Tracy, J., and Olawsky, C., 2010. Prairie Wetland Complexes as Landscape Functional Units in a Changing Climate. *BioScience* 60, 128-140.
- Johnson, W.C., Boettcher, S.E., Poiani, K.A., Guntenspergen, G., 2004. Influence of weather extremes on the water levels of glaciated prairie wetlands. *Wetlands* 24 (2), 385–398.
- Jolly, I.D., McEwan, K.L., Holland, K.L., 2008. A review of groundwater-surface water interactions in arid/semi-arid wetlands and the consequences of salinity for wetland ecology. *Ecohydrology* 1, 43-58.
- Jones, R.D., Varley, J.D., Gresswell, R.E., Jennings, D.E., Rubrecht, S.M., 1978. Fishery and aquatic management program in Yellowstone National Park, Wyoming. U.S. Fish and Wildlife Service Technical Report for 1978. Yellowstone National Park, WY.
- Junk, W.J., Fernandez Piedade, M.T., Schoengart, J., Cohn-Haft, M., Adeney, J.M., Wittmann, F., 2011. A Classification of Major Naturally-Occurring Amazonian Lowland Wetlands. *Wetlands* 31 (4), 623-640.
- van der Kamp, G., Hayashi, M., 2009. Groundwater-wetland ecosystem interaction in the semiarid glaciated plains of North America. *Hydrogeology Journal* 17 (1), 203-214.

- Klein, E., Berg, E.E., Dial, R., 2005. Wetland drying and succession across the Kenai Peninsula Lowlands, south-central Alaska. *Canadian Journal of Forest Research* 35 (8), 1931–1941.
- LaBaugh, J.W., 1987. Hydrology and chemistry of selected prairie wetlands in the Cottonwood Lake area, Stutsman County, North Dakota, 1979-82. U.S. Geological Survey Professional Paper 1431. Washington, D.C.
- LaBaugh, J.W., Winter, T.C., Rosenberry, D.O., 1998. Hydrologic Functions of Prairie Wetlands. *Great Plains Research* 8 (1), 17-37.
- Laize, C.L.R., Hannah, D.M., 2010. Modification of climate-river flow associations by basin properties. *Journal of Hydrology* 389 (1-2), 186-204.
- Lemly, J.M., Cooper, D.J., 2011. Multiscale factors control community and species distribution in mountain peatlands. *Botany* 89, 689-713.
- Long, K., Nestler, J., 1996. Hydroperiod changes as clues to impacts on Cache River Riparian Wetlands. *Wetlands* 16 (3), 379-396.
- Marsh, J.E., Timoney, K.P., 2005. How long must northern saxicolous lichens be immersed to form a waterbody trimline? *Wetlands* 25, 495-499.
- McCune, B., Mefford, M.J., 2006. PC-ORD. MjM Software, Gleneden Beach, OR.
- McMenamin, S.K., Hadly, E.A., Wright, C.K., 2008. Climatic change and wetland desiccation cause amphibian decline in Yellowstone National Park. *Proceedings of the National Academy of Sciences* 105 (44), 16988-16993.
- Merkey, D.H., 2006. Characterization of wetland hydrodynamics using HGM and subclassification methods in southeastern Michigan, USA. *Wetlands* 26 (2), 358–367.
- Mitsch, W.J., Gosselink, J.G., 2000. *Wetlands*. John Wiley & Sons, New York.
- National Climatic Data Center, 2011. NOAA. <http://www.ncdc.noaa.gov>.
- Niemuth, N.D., Wangler, B., Reynolds, R.E., 2010. Spatial and Temporal Variation in Wet Area of Wetlands in the Prairie Pothole Region of North Dakota and South Dakota. *Wetlands* 30, 1053-1064.
- Pegram, GC, Pegram, GGS, 1993. Integration of rainfall via multiquadric surfaces over polygons. *Journal of Hydraulic Engineering* 119 (2), 151-163.
- Pierce, K.L., 1979. History and dynamics of glaciation in the northern Yellowstone National Park area. U.S. Geological Survey Professional Paper 729-F. Washington, D.C.
- Pilon, P., Yue, S., 2002. Detecting climate-related trends in streamflow data. *Water Science and Technology* 45 (8), 89-104.

- Pochop, L.O., Borrelli, K., Hasfurther, V., 1985. Design characteristics for evaporation ponds in Wyoming. Final Report submitted to the Wyoming Water Research Center, Laramie, WY.
- Poff, N., Brinson, M.M., Day, J.W., 2002. Aquatic ecosystems and global climate change. Pew Center on Global Climate Change, Arlington, VA. 44 p.
- Poiani, K., Johnson, W.C., Swanson, G., Winter, T.C., 1996. Climate change and northern prairie wetlands: Simulations of long-term dynamics. *Limnology and Oceanography* 41 (5), 871-881.
- Proffitt, K.M., McEneaney, T.P., White, P.J., Garrott, R.A., 2010. Productivity and Fledging Success of Trumpeter Swans in Yellowstone National Park, 1987–2007. *Waterbirds* 33, 341-348.
- Pyke, C., 2004. Simulating vernal pool hydrologic regimes for two locations in California, USA. *Ecological Modelling* 173 (2-3), 109-127.
- R Foundation for Statistical Computing, 2009. R: A language and environment for statistical computing, Vienna, Austria.
- Rains, M.C., 2011. Water Sources and Hydrodynamics of Closed-Basin Depressions, Cook Inlet Region, Alaska. *Wetlands* 31, 377-387.
- Rawls, W., Brakensiek, D., Saxton, K., 1982. Estimation of soil-water properties: *Transactions of the ASAE* 25 (5), 1316-1328.
- Roberts, H.A., 1981. Seed banks in soils, in: *Advances in Applied Biology*. Academic Press, Cambridge, 55 p.
- Rosenberry, D., Winter, T.C., 1997. Dynamics of water-table fluctuations in an upland between two prairie-pothole wetlands in North Dakota. *Journal of Hydrology* 191 (1-4), 266-289.
- SAS Institute Inc., 2010. SAS Enterprise Guide, Cary, North Carolina.
- Schot, P., Winter, T., 2006. Groundwater-surface water interactions in wetlands for integrated water resources management - Preface. *Journal of Hydrology* 320 (3-4), 261-263.
- Shaffer, P.W., Cole, C.A., Kentula, M.E., Brooks, R.P., 2000. Effects of measurement frequency on water-level summary statistics. *Wetlands* 20, 148-161.
- Smith, L., Sheng, Y., MacDonald, G., Hinzman, L., 2005. Disappearing Arctic lakes. *Science* 308 (5727), 1429-1429.
- Smith, L.M., 2003. *Playas of the Great Plains*. University of Texas Press, Austin, TX.
- Sorenson, L., Goldberg, R., Root, T., Anderson, M.G., 1998. Potential effects of global warming on waterfowl populations breeding in the Northern Great Plains. *Climatic Change* 40 (2), 343-369.

- Stewart, I., Cayan, D., Dettinger, M., 2004. Changes in snowmelt runoff timing in western North America under a “business as usual” climate change scenario. *Climatic Change* 62 (1-3), 217-232.
- Stewart, I., Cayan, D., Dettinger, M., 2005. Changes toward earlier streamflow timing across western North America. *Journal of Climate* 18 (8), 1136-1155.
- Therneau, T.M., Atkinson, B., 2009. R port by Brian Ripley. rpart: Recursive Partitioning. R package version 3.1-45.
- Timoney, K.P., Marsh, J., 2004. Lichen trimlines in northern Alberta: establishment, growth rates, and historic water levels. *The Bryologist* 107 (4), 429–440.
- Tiner, R.W., 2003. Geographically isolated wetlands of the United States. *Wetlands* 23 (3), 494–516.
- Todd, A., Buttle, J., Taylor, C., 2006. Hydrologic dynamics and linkages in a wetland-dominated basin. *Journal of Hydrology* 319 (1-4), 15–35.
- van Tongeren, O.F.R., 1987. Cluster analysis, in: *Data Analysis in Community and Landscape Ecology*, Cambridge University Press, Cambridge, 174-203 p.
- Tsai, J.S., Venne, L.S., McMurry, S.T., Smith, L.M., 2007. Influences of land use and wetland characteristics on water loss rates and hydroperiods of playas in the Southern High Plains, USA. *Wetlands* 27 (3), 683–692.
- Upton, K.A., Jackson, C.R., 2011. Simulation of the spatio-temporal extent of groundwater flooding using statistical methods of hydrograph classification and lumped parameter models. *Hydrological Processes* 25, 1949-1963.
- United States Army Corps of Engineers, 2010. Regional supplement to the Corps of Engineers wetland delineation manual: Western mountains, valleys, and coast region (version 2.0). eds. Wakeley, J.S., Lichvar, R.W., Noble, C.V. ERDC/EL TR-10-3. US Army Engineer Research and Development Center, Vicksburg, MS.
- USDA, NRCS, 2011. The PLANTS Database. <http://plants.usda.gov>. National Plant Data Team, Greensboro, NC.
- Vega, M.R., Sierra, J.N., 1970. Population of weed seeds in a lowland rice field. *Philipp. Agricut.* 54, 1-7.
- Wallen, R., 2010. Abundance and Distribution of Yellowstone Bison, July 2010. Yellowstone Center for Resources, Yellowstone National Park, WY.
- Westerling, A.L., Turner, M.G., Smithwick, E.A.H., Romme, W.H., Ryan, M.G., 2011. Continued warming could transform Greater Yellowstone fire regimes by mid-21st century. *Proceedings of the National Academy of Sciences* 108, 13165-13170.

- Williamson, C.E., Dodds, W., Kratz, T.K., Palmer, M.A., 2008. Lakes and streams as sentinels of environmental change in terrestrial and atmospheric processes. *Frontiers in Ecology and the Environment* 6, 247-254.
- Wilmers, C.C., Getz, W.M., 2005. Gray Wolves as Climate Change Buffers in Yellowstone. *PLoS Biology* 3, e92.
- Winter, T.C., 2000. The vulnerability of wetlands to climate change: a hydrologic landscape perspective. *Journal of the American Water Resources Association* 36 (2), 305–311.
- Winter, T.C., 1986. Effect of groundwater recharge on configuration of the water-table beneath sand dunes and on seepage in lakes in the Sandhills of Nebraska, USA. *Journal of Hydrology* 86 (3-4), 221-237.
- Winter, T.C., 1999. Relation of streams, lakes, and wetlands to groundwater flow systems. *Hydrogeology Journal* 7, 28-45.
- Winter, T.C., Rosenberry, D.O., 1995. The interaction of ground-water with prairie pothole wetlands in the Cottonwood Lake area, east-central North Dakota, 1979-1990. *Wetlands* 15 (3), 193-211.
- Woo, M.K., Rowsell, R.D., 1993. Hydrology of a prairie slough. *Journal of Hydrology* 146, 175–207.
- Wurster, F.C., Cooper, D.J., Sanford, W.E., 2003. Stream/aquifer interactions at Great Sand Dunes National Monument, Colorado: influences on interdunal wetland disappearance. *Journal of Hydrology* 271 (1-4), 77–100.
- Wyman, T., 2011. 2010-2011 Annual Winter Trend Count of Northern Yellowstone Elk: Yellowstone Center for Resources, Yellowstone National Park, WY.
- Wyoming Climate Atlas, 2008, Evaporation.
<http://www.wrds.uwyo.edu/sco/climateatlas/evaporation.html>.
- Yellowstone Center for Resources, 2009. Yellowstone soils GIS layer, Yellowstone National Park, WY.
- Yellowstone National Park, 1997. Yellowstone’s Northern Range: complexity & change in a wildland ecosystem. National Park Service, Mammoth Hot Springs, WY.
- Zedler, P., 2003. Vernal pools and the concept of “isolated wetlands”. *Wetlands* 23 (3), 597-607.

9. APPENDICES

APPENDIX A. Addendum to 2009-2010 Weather Report

Methods

Annual Summaries: Weather data are publicly available since 1931 for two Northern Range weather stations within the study area, Mammoth and Tower (Fig. 1, NCDC 2011). Water year 2009 and 2010 values for temperature, total precipitation, and snow were calculated from summing (total precipitation and snow) or averaging (temperature) monthly values. 2009 and 2010 values were compared to mean values since 1931, the first year in which both Mammoth and Tower have continuous weather data. To calculate weather values, I used all months with published monthly data to determine monthly and annual averages. In the two months of 2009 (June and December) where Tower lacked data for both total precipitation and total snowfall, I used a linear regression record extension technique calculated from Mammoth data to estimate values.

Field Season Precipitation: I analyzed daily precipitation records for both Mammoth and Tower for May through August of 2009 and 2010. Additionally, on May 16 2010 I installed a HOBO tipping bucket rain gauge in the Trumpeter Lake watershed (UTM Zone 12N 549867 E, 4973438 N). This rain gauge was installed to get a better sense of a) rain inputs at our focal study site, and b) precipitation spatial variability.

Results

Annual Summaries: Both study years were near average temperature, but 2009 was a wetter year and 2010 a drier year (Table B1). 2009 surpassed the post-1931 average by 0.2°C, while 2010 was 0.3°C cooler than average. Total precipitation in 2009 was 97% and snowfall was 120% of average, while in 2010 total precipitation was only 83% and snowfall 53% of average.

Field Season Precipitation: Precipitation is correlated between the Mammoth and Tower weather stations (Pearson’s R correlation coefficient = 0.72). The largest rain event recorded was 24 mm at Mammoth on 6 August 2009. Summer precipitation most frequently falls in the form of convective summer thunderstorms, which can deliver isolated rain. Between May 13 and Aug 19, 2010 the three Northern Range precipitation gauges recorded between 105 mm (Mammoth) and 123 mm (Tower). 48% of this precipitation fell between May 28 and June 10. A water table increase was detected at wetlands after rain events (Fig. 3). The tipping bucket rain gauge and the Tower weather station each recorded ≥ 1 mm rain event on the same day 19 times over the 96 day recording period. Additionally, the tipping bucket reached ≥ 1 mm 7 times where Tower did not, and Tower recorded ≥ 1 mm 8 times where the tipping bucket did not.

Table A1. Temperature and precipitation deviation from historic average, 2009 and 2010. Mam = Mammoth, Tow = Tower, NR = Northern Range value, the mean of Mammoth and Tower.

	2009	2010
Mam Temp (°C)	+0.2	-0.4
Tow Temp (°C)	+0.2	-0.2
NR Temp (°C)	+0.2	-0.3
Mam Ppt	97%	81%
Tow Ppt	97%	85%
NR Ppt	97%	83%
Mam Snow	111%	45%
Tow Snow	129%	60%
NR Snow	120%	53%

APPENDIX B. Location of all wells

Table B1. Log of all 138 wells used in the study.

Well #	Wetland	Easting	Northing	Elev (m)	Well Type
1	DC	543566	4976420	2047	shape3-mag2
4	DC	543510	4976320	2047	shape3-mag2
9	DC	543616	4976400	2048	shape3-mag2
17	BM	547854	4973670	1942	shape1-mag1
21	BM	547835	4973690	1943	shape1-mag1
22	BM	547949	4973640	1943	shape1-mag1
26	TF	550143	4973550	1898	shape2-mag2
29	TF	550155	4973580	1898	shape2-mag2
32	TF	550092	4973570	1898	shape2-mag2
34	TL	549963	4973720	1897	shape2-mag2
36	TL	549993	4973780	1897	shape2-mag2
38	TL	549950	4973950	1897	shape3-mag1
40	TL	549664	4973950	1897	n/a
42	TL	549747	4973700	1897	shape3-mag1
44	BP	520591	4985781	1799	stable
45	DW	520435	4985686	1809	perched
47	RL	520381	4985529	1825	perched
48	MC	520841	4985700	1804	stable
51	MC	520834	4985770	1797	stable
53	MC	520784	4985780	1799	stable
56	BP	520594	4985770	1800	stable
57	BP	520597	4985873	1798	stable
59	DW	520413	4985700	1810	perched
63	CR	550195	4973120	1924	shape1-mag1
66	CR	550294	4973029	1924	shape1-mag1
69	CR	550305	4973068	1922	shape1-mag1
71	CR	550357	4973140	1919	shape1-mag1
74	LT	549470	4973915	1898	shape1-mag2
75	BD	546614	4974912	1910	shape1-mag1
76	SN	548471	4974500	1901	shape3-mag2
79	SN	548509	4974500	1899	shape3-mag2
81	SN	548466	4974540	1900	shape3-mag2
85	SN	548505	4974560	1903	shape3-mag2
92	DW	520460	4985610	1813	Perched
93	DW	520458	4985620	1812	Perched
96	DW	520449	4985650	1810	shape2-mag2
100	DW	520451	4985700	1808	Perched
102	DW	520441	4985760	1806	Stable
105	RL	520428	4985370	1826	Stable
107	RL	520359	4985470	1825	shape3-mag2
110	CP	520821	4971730	2284	n/a
112	CP	520764	4971670	2284	n/a

114	CP	520820	4971712	2284	n/a
115	SV	552100	4974480	1924	shape1-mag1
118	SV	552055	4974550	1923	shape1-mag1
121	SV	552070	4974460	1924	shape1-mag1
123	IP	551940	4974190	1925	shape1-mag2
125	IP	551941	4974190	1925	shape1-mag2
127	IP	551978	4974125	1925	shape2-mag2
128	IP	552033	4974130	1925	shape1-mag2
130	BW	521983	4974430	2270	shape1-mag1
133	BW	521972	4974280	2269	shape1-mag1
136	BW	522072	4974220	2269	n/a
137	BE	522332	4973990	2260	shape2-mag2
140	BE	522574	4973810	2260	shape2-mag2
142	BE	522338	4973860	2260	shape2-mag2
144	BE	522409	4973956	2260	shape2-mag2
148	FF	532455	4978196	2054	n/a
152	RP	524225	4978480	1924	shape3-mag2
155	RP	524176	4978560	1923	shape3-mag2
158	RP	524135	4978540	1925	shape3-mag2
161	BD	546619	4974870	1911	shape1-mag1
164	BD	546578	4974890	1911	shape1-mag1
168	BD	546619	4974920	1910	shape1-mag1
170	BD	546615	4974900	1910	shape1-mag1
172	BM	547937	4973570	1941	n/a
174	SG	534905	4979120	2089	shape1-mag2
177	SG	534765	4979340	2080	n/a
180	SG	534880	4979270	2086	shape2-mag2
182	SG	534835	4979260	2086	shape2-mag2
184	MB	524875	4978880	1861	shape3-mag1
187	MB	524953	4978980	1853	shape3-mag1
189	MB	524943	4978880	1859	shape3-mag1
192	MB	524926	4978910	1858	shape3-mag1
195	WA	551200	4973610	1924	n/a
197	WA	551448	4973590	1921	stable
200	WA	551361	4973420	1921	stable
203	LT	549468	4973860	1899	shape2-mag2
210	LT	549305	4974000	1899	shape3-mag1
213	LT	549480	4974030	1896	shape2-mag2
217	US	523563	4983270	1786	shape3-mag1
220	US	523581	4983270	1785	shape3-mag1
223	US	523595	4983300	1783	shape3-mag1
226	US	523596	4983280	1784	shape3-mag1
229	OR	523265	4983550	1825	shape2-mag2
231	OR	523288	4983600	1826	shape2-mag2
233	OR	523348	4983630	1823	shape2-mag2
236	OR	523331	4983580	1824	shape2-mag2
239	FF	532336	4978330	2054	n/a

242	FF	532293	4978280	2053	n/a
243	BG	545994	4974950	1945	shape1-mag1
246	BG	545957	4974960	1944	shape1-mag1
248	BG	545969	4974930	1945	shape1-mag1
251	BG	545945	4974940	1944	n/a
254	BG	545950	4974946	1944	n/a
255	CP	520805	4971670	2284	shape1-mag1
257	BW	522122	4974300	2271	n/a
258	RP	524215	4978540	1922	shape3-mag2
260	RP	524176	4978540	1923	shape3-mag2
267	FF	532390	4978272	2054	n/a
271	LT	549386	4973860	1899	shape1-mag2
272	MC	520856	4985740	1799	stable
273	MC	520855	4985740	1799	stable
276	IP	551983	4974117	1925	shape2-mag2
279	SV	552065	4974446	1925	shape1-mag1
282	TW	549943	4972880	1946	n/a
283	TW	550079	4973150	1930	n/a
286	TW	549784	4973290	1925	n/a
288	TW	549803	4973320	1925	n/a
291	TW	549981	4972970	1936	n/a
295	FF	532324	4978310	2053	n/a
296	TL	549917	4973640	1897	shape3-mag2
299	TL	549864	4973620	1897	shape3-mag1
301	TW	549817	4973430	1915	n/a
303	BM	548011	4973580	1941	n/a
305	BM	547944	4973610	1942	shape1-mag1
307	DC	543570	4976400	2047	shape3-mag2
308	CP	520793	4971692	2284	shape1-mag1
309	BP	520589	4985815	1798	shape1-mag2
310	BW	522054	4974226	2269	shape1-mag1
311	SV	552049	4974494	1924	n/a
313	TL	549752	4973812	1897	shape2-mag2
316	TL	549897	4973857	1897	shape2-mag2
320tf	TF	550163	4973597	1900	n/a
320wa	WA	551443	4973344	1924	n/a
322	WA	551430	4973573	1920	stable
323	WA	551280	4973564	1921	shape1-mag1
326	CR	550294	4973090	1922	shape1-mag1
328	SG	534790	4979319	2081	n/a
334	BE	522325	4973975	2260	shape2-mag2
336	BG	545964	4974943	1944	shape1-mag1
338	TW	550256	4973451	1908	n/a
341	TL	549755	4973918	1897	n/a
342	TL	549721	4973768	1897	n/a
343	TL	549790	4973655	1898	n/a
344	TL	549921	4973618	1897	n/a

345	TL	549978	4973677	1897	n/a
347	SN	548498	4974538	1900	shape3-mag2
348	TW	550272	4973462	1908	n/a
349	OR	523303	4983572	1824	shape2-mag2
350	IP	552004	4974165	1925	shape2-mag2
351	TL	549888	4973651	1856	shape2-mag2
352	WA	551429	4973574	1920	shape1-mag1

APPENDIX C. Aerial photography dates

Table C1. Day, or window of days, on which air photos were taken in each year. Note that 1969 lacked photos for BD and BG so photos from 1971 were substituted.

Year	Wetlands near Mammoth	Wetlands near Tower
1954	Aug 11-18	Aug 11-18
1969	Sep 8	Sep 8
1971	-	Sep 10
1991	Jul 4	Aug 3
1994	Jun 26	Aug 24
1998	Aug 5	Aug 5
2001	Jul 1	Jul 1
2006	Sep 29	Sep 29
2009	Aug 27	Aug 27

APPENDIX D. Hydraulic conductivity measurements in the Trumpeter Lake watershed

Saturated hydraulic conductivity (K_s) was calculated via the double-ring infiltrometer method and via the Hvorslev slug test. The two methods yielded quite different results, possibly a function of depth (van der Kamp and Hayashi 2009). This was unexpected in an unconsolidated till system with poor soil development, since vertical and horizontal K_s to be roughly equivalent. Results from each location tested follow in these tables. Note that in Table G1 well 296 at 35 cm depth should equal the “TL lake seds.” at 35 cm depth since these tests were performed in the same location. The values were 0.037 and 0.26 m/day, respectively. This narrow line of evidence suggests an error of one degree of magnitude. I am not sure where the error originated.

Table D1. K_s calculated from the double-ring infiltrometer test.

Location	Depth	K_s (m/day)
Upland TW #2	Surface	0.8
Upland TW	Surface	0.87
Upland TW	50 cm	0.56
TW local depression, <i>Juncus</i> veg.	Surface	1.7
TL lake seds., well 296	Surface	0.13
TL lake seds., well 296	35 cm	0.26

Table D2. K_s calculated from the Hvorslev slug test.

Well	K_s (m/day)
341	0.011
38	0.043
342	0.025
343	0.003
299	0.016
296	0.037
344	0.0029
345	0.02
32	1.5
29	0.23
26	0.022
301	0.04

APPENDIX E. Trumpeter Lake focal site analysis




Figure E1. Trumpeter Lake surface water extent through time. A) shows lake perimeter delineated directly from air photographs. B) shows perimeter calculated by accounting for evaporation and precipitation between photo date and the standardization date, August 15. Land surface was produced by creating a TIN from a basin-wide topographic survey conducted in 2010. 2010 lake perimeter was calculated from field data directly in absence of an aerial photograph.

Table E1. Trumpeter Lake size derived from aerial photos. Volumes were calculated using the fill function in a TIN layer in ArcGIS. Since lake size changes throughout the summer and aerial photos were taken on different days in different years, I estimated lake size for the standardized date of August 15 by adjusting for the precipitation recorded at the Tower rain gauge (NCDC 2011) and the estimated lake evaporation (Pochop et al. 1985, WCA 2008)

Photo Date	Photo Date		15 Aug Standardization	
	Area (m²)	Volume (m³)	Area (m²)	Volume (m³)
11-18 Aug 1954	106,000	168,000	105,000	168,000
8 Sep 1969	126,000	249,000	128,000	249,000
3 Aug 1991	71,000	66,000	69,000	63,000
24 Aug 1994	68,000	61,000	69,000	63,000
5 Aug 1998	100,000	152,000	99,000	147,000
1 Jul 2001	72,000	69,000	67,000	57,000
29 Sep 2006	35,000	14,000	30,000	10,000
27 Aug 2009	48,000	27,000	50,000	28,000
xx Xxx 2010	n/a	n/a	35,000	14,000

APPENDIX F. Soil seed bank study species list

Table F1. Plant species germinating in the soil seed bank study, listed with their wetland indicator status. n = number of wetlands, out of 12, from whose plots a species germinated. Nomenclature follows USDA PLANTS (2011).

Species Name	Wetland Indicator	
	Status	n
<i>Achillea millefolium</i> var. <i>lanulosa</i>	FACU	1
<i>Agrostis scabra</i>	FAC	2
<i>Agrostis stolonifera</i>	FAC	3
<i>Alyssum desertorum</i>	N/A	4
<i>Androsace septentrionalis</i>	FAC-	3
<i>Antennaria microphylla</i>	N/A	1
<i>Artemisia frigida</i>	N/A	1
<i>Astragalus agrestis</i>	FACW-	2
<i>Bromus japonicus</i>	UPL	6
<i>Bromus tectorum</i>	N/A	1
<i>Camelina microcarpa</i>	N/A	1
<i>Cerastium fontanum</i>	FACU	1
<i>Ceratocephala testiculata</i>	N/A	3
<i>Cirsium arvense</i>	FACU+	2
<i>Collinsia parviflora</i>	N/A	2
<i>Collomia linearis</i>	FACU	2
<i>Corydalis aurea</i>	N/A	1
<i>Cryptantha ambigua</i>	N/A	1
<i>Dasiphora fruticosa</i>	FAC-	1
<i>Descurainia pinnata</i> var. <i>nelsonii</i>	N/A	1
<i>Descurainia sophia</i>	N/A	2
<i>Draba nemorosa</i>	N/A	11
<i>Epilobium ciliatum</i>	FACW-	11
<i>Erysimum inconspicuum</i>	N/A	2
<i>Festuca idahoensis</i>	FACU*	1
<i>Fragaria virginiana</i>	FACU*	3
<i>Gnaphalium palustre</i>	FAC+	3
<i>Juncus arcticus</i>	FACW+	3
<i>Juncus ensifolius</i>	FACW	2
<i>Juncus longistylis</i>	FACW	1
<i>Juncus nodosus</i>	OBL	1
<i>Juncus torreyi</i>	FACW	1
<i>Melilotus officinalis</i>	FACU	1
<i>Mentha arvensis</i>	FACW-	3
<i>Monolepis nuttalliana</i>	FAC-	3
<i>Myosurus minimus</i>	OBL	2

<i>Poa palustris</i>	FAC	12
<i>Puccinellia nuttalliana</i>	FACW+	1
<i>Ranunculus macounii</i>	OBL	3
<i>Ranunculus sceleratus</i>	OBL	7
<i>Rumex maritimus</i>	FACW+	6
<i>Taraxacum officinale</i>	FACU	7
<i>Thlaspi arvense</i>	N/A	5
<i>Trifolium hybridum</i>	FAC	1
<i>Typha latifolia</i>	OBL	2
<i>Veronica peregrina</i>	OBL	5

APPENDIX G. Soil seed bank study plot stratification procedure

I explored multiple ways to analyze soil seed bank data. As explained in the text, I resolved on a classification stratifying plot classes by their relative elevation from the lowest plot (0%) up to the surrounding upland elevation (100%). Elevation classes were chosen according to two criteria, 1) I attempted to lump percentages in close proximity to each other (e.g. given the percentages 0, 17, 20, and 25 I grouped 0's together and 17, 20 and 25's together), and 2) I attempted to create groups of similar size ($20 \leq n \leq 24$ for six of the seven classes). Table J1 shows the raw data for number of OBL and FACW species germinating in each plot, grouped by elevation class. The total number of wetland species germinating per plot was lower than desired.

An alternative classification that was rejected used each plot's elevation above the 2009 maximum water level in a particular wetland basin (cm). The advantage of this method was that it placed all wetlands on the same absolute scale, and elevation classes were described in centimeters rather than as percentages. This method incorporated the ecological importance of the seasonal high water level. However, the drawback to this classification was that it tended to decrease the diversity of wetlands represented in the plot elevation classes (e.g. only 7 of 12 wetlands were represented in the lowest class; Trumpeter Lake provided 8 of the 24 plots in the highest class). This classification revealed no strong trends in the distribution of germinating wetland species. In fact, wetland species starkly peaked in the highest elevation class, 140-342 cm above 2009 peak. Plots in this class were dominated by Trumpeter Lake plots, a particularly productive wetland. Thus, a high percentage of plots from one wetland and no pattern outside of this elevation class discounted my trust in this analysis.

Table G1. Raw data showing number of wetland species germinating per plot in the soil seed bank study. Plots are grouped in columns according to their relative wetland elevation, represented by percentage from wetland bottom to upland vegetation.



	0%	17-25%	33-40%	50%	60-67%	75-83%	100%
	1	0	1	2	0	1	0
	0	0	0	2	2	1	0
	0	0	1	1	4	1	2
	1	1	1	0	1	0	1
	1	1	1	3	3	1	0
	0	0	0	2	1	0	1
	0	0	2	0	2	2	1
	0	0	1	0	1	1	1
	0	1	0		1	1	1
	0	1	1		1	1	1
	0	3	2		3	0	2
	0	0	1		1	0	0
	2	1	0		0	1	1
	1	1	0		0	1	0
	1	1	0		2	4	2
	0	0	0		2	0	4
	0	1	1		2	0	1
	2	0	0		0	0	0
	3	2	3		3	2	0
	0	1	2		2	2	0
	0	2				2	1
		1				0	1
		1				5	1
		0				0	0
mean	0.6	0.8	0.9	1.3	1.6	1.1	0.9
SE	0.19	0.16	0.20	0.41	0.26	0.26	0.19

Table G2. Mann Whitney U test statistics from comparing each elevation class to all others. The top right triangle is the test z-score, while the bottom left triangle shows the associated p-values. Asterisks highlight significance at $p < 0.05$. See Fig. 12 for bar graph.


Elev. Class	0%	17-25%	33-40%	50%	60-67%	75-83%	100%
0%		-1.04	1.23	1.51	2.88*	-1.57	-1.36
17-25%	0.3		0.33	1.09	2.45*	-0.7	-0.38
33-40%	0.23	0.74		0.83	-2*	-0.34	0
50%	0.14	0.28	0.41		-0.53	0.57	0.89
60-67%	0.01*	0.02*	0.05*	0.6		1.64	2.13*
75-83%	0.12	0.48	0.74	0.57	0.11		0.35
100%	0.18	0.7	1	0.39	0.04*	0.73	





APPENDIX H. Example wetland hydrographs from the seven classes

Figure H1. Illustrated are hydrographs of the seven wetland classes derived from well hydrograph analysis. Each pane is an example wetland from a given class. Summer hydrographs for all wells at a wetland are illustrated for both years. A given well is illustrated with the same color in each year. 2009 hydrographs are solid lines and 2010's are dashed. Black lines are staff gauges.



70





APPENDIX I. Yellowstone River annual discharge




Figure I1. Mean annual discharge (Q) on the Yellowstone River at the Corwin Springs gauging station. Dots represent annual mean Q's for 1911-2010, black line represents the 10-year running average, and gray line is the mean. Data revealed an oscillating pattern of wetter and drier years, but no comprehensive trajectory over the past century. This figure's trends were extended back to the early 18th century by Graumlich et al. (2003). Their data indicated that the wettest period was circa 1900 and the longest prolonged dry spell was 1780-1870.

APPENDIX G. Photographs of one wetland from each class



Figure G1. Stable surface water class, wetland RL.



Figure G2. Perched water table class, wetland BP.



Figure G3. Seasonally and yearly variable water table class, wetland BD.



Figure G4. Peat soils with recovering water tables class, wetland DC.



Figure G5. Mineral soils with recovering water tables class, wetland MB.



Figure G6. Interannually variable water table persistence, wetland OR.



Figure G7. Spatially variable water table patterns, wetland TL.